

V.M. RED'KOV

On solutions of Schrödinger and Dirac equations in spaces of constant curvature, spherical and elliptical models

Exact solutions of the Schrödinger and Dirac equations in generalized cylindrical coordinates of the 3-dimensional space of positive constant curvature, spherical model, have been obtained. It is shown that all basis Schrödinger's and Dirac's wave functions are finite, single-valued, and continuous everywhere in spherical space model S_3 . The used coordinates (ρ, ϕ, z) are simply referred to Euler's angle variables (α, β, γ) , parameters on the unitary group $SU(2)$, which permits to express the constructed wave solutions $\Psi(\rho, \phi, z)$ in terms of Wigner's functions $D_{mm'}^j(\alpha, \beta, \gamma)$. Specification of the analysis to the case of elliptic, $SO(3.R)$ group space, model has been done. In so doing, the results substantially depend upon the spin of the particle. In scalar case, the part of the Schrödinger wave solutions must be excluded by continuity considerations, remaining functions are continuous everywhere in the elliptical 3-space. The latter is in agreement with the known statement: the Wigner functions $D_{mm'}^j(\alpha, \beta, \gamma)$ at $j = 0, 1, 2, \dots$ make up a correct basis in $SO(3.R)$ group space. For the fermion case, it is shown that no Dirac solutions, continuous everywhere in elliptical space, do exist. Description of the Dirac particle in elliptical space of positive constant curvature cannot be correctly in the sense of continuity adjusted with its topological structure.

Key words: spherical, elliptical, geometry, Schrödinger and Dirac equation, Wigner functions, Euler angles, continuity, curvature, spin, topology.

1. Introduction

The main quantum mechanical equations (say, for definiteness, scalar Schrödinger and fermion Dirac ones) are assumed to have the corresponding wave functions be given in the whole space. So, the global structure of the space model, its topological properties, may be of substantial significance in the quantum mechanical context.

In the present paper, two canonical examples, Schrödinger's and Dirac's equations on the manifolds of the unitary $SU(2)$ and orthogonal $SO(3.R)$ groups have been examined in detail. These manifolds provide us with topologically different variants of the most simple Riemannian space, the 3-space of constant positive curvature, spherical S_3 and elliptical $\tilde{S}_3$ models respectively. These two groups, $SU(2)$ and $SO(3.R)$, are widely used in many applications, in the same time the groups give simple and familiar realizations of these two geometrical models, that can be followed in detail by quite elementary mathematical tools. So, in these simple systems, the influence of the topology in quantum mechanics can be traced in terms of exact solutions.

In the paper, solutions of the main equations in the space S_3 and $\tilde{S}_3$ will be constructed in the coordinates (ρ, ϕ, z) which can be considered as extension of the familiar cylindrical coordinates in flat space. These curved coordinates (ρ, ϕ, z) can be referred to the Euler angle variables: $\alpha = \phi + z$, $\beta = 2\rho$, $\gamma = \phi - z$. The latter gives possibility to express any given wave solution (Schrödinger's or Dirac's) in terms of Wigner's functions $D_{mm'}^j(\alpha, \beta, \gamma)$.

There exists one other point making the quantum mechanical problem under consideration of interest in connection with the old and eminent theorem in the theory of compact groups. The theorem by Peter and Weyl ([1], see its formulation in [2] states:

Teorema I: Let $\hat{G} = \{T^s\}$ be the set of all irreducible unitary representations $\sqrt{d_s} D_{jk}^s(g)$, $s \in \hat{G}$, $1 \leq j, k \leq d_s$, d_s stands for the dimension of the representation. Matrix elements D_{jk}^s of the representation T^s make up the full orthogonal basis in $L^2(G)$. For any function $U(g) \in L^2(G)$ there exists decomposition

$$U(g) = \sum_{s \in \hat{G}} \sum_{j,k=1}^s C_{jk}^s D_{jk}^s(g), \quad \int_G |U(g)|^2 dg = \sum_{s \in \hat{G}} d_s \sum_{j,k=1}^{d_s} |C_{jk}^s|^2.$$

The Theorem I can be significantly strengthened.

Teorema II: Let $f(g)$ be a function continuous in G . For any $\epsilon > 0$ there exists a linear combination of matrix elements such that

$$|f(g) - \sum_{s=1}^{N_\epsilon} [\sum_{j,k=1}^{d_s} C_{jk}^s D_{jk}^s(g)]| \leq \epsilon \quad g \in G.$$

This Theorem II states that any function $f(g)$ continuous in G can be represented by the series in terms of $D_{jk}^s(g)$, and this series coincides with the initial function $g(g)$ in all the points of G . By the Theorem I, the function $f(g)$, discontinuous somewhere in G , can be exactly approximated by the series only at the points of continuity.

It should be noted that in spite of the wide use the Euler's parameters in the theory of groups $SU(2)$ and $SO(3.R)$ in the literature there exist full confusion in pointing out a domain in which the variables (α, β, γ) should change to be correct in context of the Peter-Weyl theorem II – see [3-20]. In that sense, the most interesting analysis of the problem had been given in [21].

As a rule, they operate only with the explicit form of matrix elements $D_{jk}^s(\alpha, \beta, \gamma)$ and with the orthogonality conditions for them. At this any clear distinction between different variants $G(\alpha, \beta, \gamma)$ has not been achieved. In the literature we can see each of the following domains:

$$G = 2\pi \otimes \pi \otimes \pi, \quad G = \pi \otimes \pi \otimes 2\pi, \quad G = 2\pi \otimes \pi \otimes 2\pi, \\ G = 4\pi \otimes \pi \otimes 4\pi, \quad G = 2\pi \otimes \pi \otimes 4\pi, \quad G = 4\pi \otimes \pi \otimes 2\pi, \quad G = 2\pi \otimes 2\pi \otimes 2\pi.$$

As a rule, without any special examining, it is tacitly assumed that all functions involved are continuous. In the essence, two different concepts, a function continuous in the manifold $SU(2)$ or $SO(3.R)$, and a function continuous with respect to three variables (α, β, γ) , are not distinguished. Till now, no special treatment joining different points of view does not exist. So other additional aim of the present work is to give such an analysis that might give possibility to use the Peter-Weyl theorem II correctly. In the same time, the question of parametrization of unitary and orthogonal groups by Euler variables must be fully clarified in the framework of the main goal of the present work – to find exact solutions of Schrödinger and Dirac equations in the spaces of constant positive curvature. Because it is evident in advance that the main requirements for these are their finiteness and continuity in all the points of spherical S_3 and elliptic $\tilde{S}_3$ spaces.

In the beginning (§1 - §2) we are consider solutions of the scalar Schrödinger equation in the spherical space

$$n_0^2 + n_1^2 + n_1^2 + n_1^2 = 1, \quad n_a = n_a(\rho, \phi, z),$$

in the the generalized cylindrical coordinates

$$n_0 = \cos \rho \cos z, \quad n_3 = \cos \rho \sin z, \quad n_1 = \sin \rho \cos \phi, \quad n_2 = \sin \rho \sin \phi,$$

$$G(\rho, \phi, z) : \quad \rho \in [0, \pi/2], \quad \phi, z \in [-\pi, +\pi]$$

That cylindrical parametrization has some peculiarities. The domain G may be divided into three parts: G_1, G_2, G_3 dependently on the values of ρ : $\rho \in (0, \pi/2), \rho = 0, \rho = \pi/2$. At this the domains $G_2(0, \phi, z)$ and $G_3(\pi/2, \phi, z)$ correspond to 1-dimensional closed curves:

$$G_2 : \quad n_0 = \cos z, \quad n_3 = \sin z, \quad n_0^2 + n_3^2 = 1,$$

$$G_3 : \quad n_1 = \cos \phi, \quad n_2 = \sin \phi, \quad n_1^2 + n_2^2 = 1.$$

Therefore, any wave function $\Psi(\rho, \phi, z)$ continuous in the space S_3 must obey a number of definite restrictions:

$$(G_1) : \quad \Psi(0, \rho < \pi/2, \phi, z) = \Psi(0, \rho < \pi/2, \phi \pm 2\pi, z \pm 2\pi),$$

$$(G_2) : \quad \Psi(0, \phi, z) = \{ \Psi(z) = \Psi(z \pm 2\pi), \text{ or } \text{const} \},$$

$$(G_3) : \quad \Psi(0, \phi, z) = \{ \Psi(\phi) = \Psi(\phi \pm 2\pi), \text{ or } \text{const} \}.$$

Taking the elliptical model (§3) will much change the all treatment. As "Cartesian" may be used the known Gibbs [22] coordinates $\vec{c} = (c_1, c_2, c_3)$ ¹


$$O(\vec{c}) = I + 2 \frac{\vec{c}^\times + (\vec{c}^\times)^2}{1 + \vec{c}^2};$$

as seen to any pair of inversely directed vectors of infinite length corresponds the same matrix, point in elliptic space:

$$\vec{c}^+ = +\infty \vec{c}_0, \quad \vec{c}^- = -\infty \vec{c}_0, \quad \vec{c}_0^2 = 1, \quad O(\vec{c}^+) = O(\vec{c}^-) = I + 2(\vec{c}_0^\times)^2.$$

Cylindrical parametrization of the space $\tilde{S}_3$ has its own peculiarities. The domain $\tilde{G}(\rho, \phi, z)$ should be decrease twice as much and new identification rule for the boundary points should be accepted (here only one way to chose the domain $\tilde{G}$ is detailed):

Identification in $\tilde{G}_1^{(+1)}(\phi, z) : \rho \neq 0, \pi/2$



$$\tilde{G}_2(0, \phi, z) : \quad \rho = 0, \quad \vec{c} = (0, 0, \tan z), \quad \phi - \text{нeмaя};$$

$$\tilde{G}_3(0, \phi, z) : \quad \rho = \pi/2, \quad \vec{c} = \infty (\cos \phi, \sin \phi, 0), \quad z - \text{мyтe}.$$

The requirement of continuity in elliptic space $\tilde{S}_3$ must be agreed with these identification rules. In that way it is shown that the obtained solutions of Schrödinger equation in S_3 can

¹For detailed and systematic exposition of the theory of the rotation group in terms of $\vec{c}$ -parameters and much further extensions see the book [10].

be divided into two classes as follows: the first type functions turn out to be discontinuous in space $\tilde{S}_3$ and therefore they should be excluded from the list of "good" basis functions. The second type solutions are continuous both in spherical and elliptic spaces and they make up the full basis system in $\tilde{S}_3$. In view of referring the coordinates (ρ, ϕ, z) to Euler variables $SU(2)$ и $S0(3.R)$, the above can be formulated (§4 -§5) as follows: basis for continuous functions on the group $SU(2)$ consists of $D_{m\sigma}^j(\alpha, \beta, \gamma)$ with $j = 0, 1/2, 2, 3/2, \dots$; basis of continuous functions on the group $S0(3)$ consists of $D_{m\sigma}^j(\alpha, \beta, \gamma)$ with $j = 0, 1, 2, \dots$.

It is natural to examine in analogous way the case of Dirac particle in spherical and elliptical spaces. In §6 the Dirac equation is specified in spherical space; cylindrical coordinates and tetrad are used. The wave solutions $\Psi_{\epsilon, \lambda, m, k}(t, \rho, \phi, z)$, eigenfunction of four operators have been looked for. The problem is reduced to a linear system of radial equations for two functions $f_1(\rho)$, $f_2(\rho)$. In §7 a special transformation is found $f_1(\rho)$, $f_2(\rho) \Rightarrow G_1(\rho)$, $G_2(\rho)$, which permits to solve the problem for $G_1(\rho)$, $G_2(\rho)$ in terms of hypergeometric functions. Further (§8- §11) the quantization rules for ϵ, λ, m, k are to be derived and corresponding wave functions are to be constructed.

The main requirement is: these wave functions must be finite and continuous in the space S_3 . The formulation of the latter condition in the curved space is much more complex in case of fermion. The matter is that besides the simple technical complication – the number of components of a bispinor is four – the mathematical formalism itself, by Tetrode-Weil-Fock-Ivanenko [23-25], involves an additional local gauge transformations – for the case under consideration those are local tetrad $SU(2)$ rotations. These gauge transforms connected with the freedom in specifying tetrads are singular so that the explicit form of continuity requirement depends upon the occasional choice of the tetrad.

Strictly speaking, the same problem is in flat space as well. Indeed, the wave functions being single-valued in Cartesian tetrad do not look single-valued in the cylindrical tetrad. Therefore, if one starts with that cylindrical tetrad one will face the problem – which solutions are continuous and which ones are not.

In the case of spherical space S_3 the all situation is more complex though principally the same. The role of a Cartesian tetrad basis in S_3 belongs to the conformally flat tetrad. In Supplement A, an explicit form of the spinor gauge matrix $B(\rho, \phi, z)$ referring cylindrical and conformally-flat tetrads is calculated. The main point is that the fermion wave function may be discontinuous in the cylindrical basis but it must become continuous in conformally-flat basis.

In particular, because the gauge matrix $B(\rho, \phi, z)$ contains the trigonometrical functions of the half angles $\phi/2$ and $z/2$ the quantum numbers m and k must take the half-integer values. In quantum number λ , the proper value of the curved helicity operator, the energy levels are twice degenerated: $\epsilon = \sqrt{M^2 + \lambda^2}$. Finally, all hypergeometric function involved become polynomials when the λ is quantized as follows: $\lambda = \pm 3/2, 5/2, 7/2, \dots$ In §11 the special analysis, showing that all the constructed fermion wave functions are continuous in all the space S_3 , is given. In addition (§9), all the solutions with $\lambda = \pm 1/2$ must be excluded by continuity reason.

In §12 the Dirac equation in elliptic space $\tilde{S}_3$ has been considered. Application of the tetrad formalism in this model has its own peculiarities. In particular, because of $(2 \rightarrow 1)$ -correspondence between points of spherical and elliptic manifolds any tetrad given in the space S_3 provides us with two different tetrads in elliptical space $\tilde{S}_3$. For instance, the conformally flat retrad after transforming to c^i -coordinates in $\tilde{S}_3$ lead us to a pair of those

$$e_{(i)j}^{(\delta)}(c) = \frac{\delta_{ij}}{\delta\sqrt{1+c^2}} - \frac{c^i c^j}{(1+c^2)(1+\delta\sqrt{1+c^2})}, \quad \delta = \pm 1 ,$$

Such a doubling is a consequence of the following: elliptic manifold can be constructed from spherical one by two ways: on the base of the half-space S_3 containing the point $n_a = (+1, 0, 0, 0)$ or on the base of the half-space s_3 containing another point $n_a = (-1, 0, 0, 0)$.

In the same manner, cylindrical coordinates and tetrads of spherical space produce a pair of coordinate systems and tetrad in elliptic space which manifests itself as freedom to use any of two different domains: $(\phi, z) \in \tilde{G}^{(+1)}(\phi, z)$ and $(\phi, z) \in \tilde{G}^{(-1)}(\phi, z)$. We have been calculated the tetrad gauge matrix $L_{(\delta)}$ referring cartesian and cylindrical tetrads, then calculate the corresponding spinor gauge matrices (for both cases $\delta = +1, -1$):

$$L_{(\delta)}(\rho, \phi, z) e_{cart}^{(\delta)}(c) = e_{cyl}^{(\delta)}(\rho, \phi, z) , \quad \psi_{cart}^{(\delta)} = \pm B_{(\delta)}^{-1}(\rho, \phi, z) \psi_{cyl}^{(\delta)} .$$

So there exist two cartesian representations for fermion wave functions and any of these can be used in considering the requirement of continuity for fermion wave functions².

It is shown that all Dirac wave solutions constructed continuous in spherical space model become discontinuous when specifying them to elliptic model. Therefore, description of the Dirac particle in elliptical space of positive constant curvature cannot be correctly in the sense of continuity adjusted with its topological structure.

The paper is divided into three parts. In the first part (§2 -§5), we consider Schrödinger equation, in spherical and then in elliptical space models. In the second part (§6 - §11) construct exact solutions of the Dirac equation on the sphere S_3 , all the functions founded are continuous in the space. Third part (§12 -§13) treats the case of Dirac equation in elliptical space. It is shown that no Dirac solutions, continuous everywhere in elliptical space, do exist.

PART I. SCHRÖDINGER EQUATION IN SPHERICAL AND ELLIPTICAL SPACES, EILER PARAMETRIZATION AND WIGNER D -FUNCTIONS

2. Schrödinger equation in spherical space S_3 .

The cylindrical coordinates on the sphere S_3 are defined by

$$n_0 = \cos \rho \cos z , \quad n_3 = \cos \rho \sin z , \quad n_1 = \sin \rho \cos \phi , \quad n_2 = \sin \rho \sin \phi ,$$

$$dl^2 = [d\rho^2 + \sin^2 \rho d\phi^2 + \cos^2 \rho dz^2] , \quad \rho \in [0, \pi/2] , \quad \phi, z \in [-\pi, \pi] . \quad (2.1)$$

For a small part of the space S_3 in vicinity of $n_a = (1, 0, 0, 0)$ the quantities (ρ, ϕ, z) become ordinary (flat) cylindrical coordinates. The Schrödinger generally covariant equation

$$H \Phi = \epsilon \Phi , \quad H = -\frac{1}{2} \frac{1}{\sqrt{g}} (\partial/\partial x^i \sqrt{g} g^{ij} \partial/\partial x^j) ,$$

where $\epsilon = E/(\hbar^2/MR^2)$ stands for the particle energy, being specified in the system (2.1) will take the form

$$[\sin^{-1} \rho \cos^{-1} \rho (\partial_\rho \sin \rho \cos \rho \partial_\rho) + \sin^{-2} \rho \partial_\phi^2 + \cos^{-2} \rho \partial_z^2 + 2\epsilon] \Phi = 0 . \quad (2.2a)$$

²Two other possibilities resulting from combination of the above two have been considered as well.

Solutions can be searched as cylindrical waves $\Phi(\rho, \phi, z) = e^{+iM\phi} e^{+iKz} R(\rho)$; for $R(\rho)$ one has the equation

$$\begin{aligned} & [d^2/d\rho^2 + (\cos \rho / \sin \rho - \sin \rho / \cos \rho) d/d\rho + \\ & + 2\epsilon - M^2 / \sin^2 \rho - K^2 / \cos^2 \rho] R(\rho) = 0 . \end{aligned} \quad (2.2b)$$

The function $R(\rho)$ reduces to hypergeometric one

$$R(\rho) = \sin^a \rho \cos^b \rho F(A, B, C; \cos^2 \rho) , \quad (2.2)$$

$$A = (a + b + 1 - \sqrt{2\epsilon + 1})/2 , \quad B = (a + b + 1 + \sqrt{2\epsilon + 1})/2 , \quad C = (b + 1) .$$

The wave function $\Phi(\rho, \phi, z)$ will be single-valued in S_3 (in the part G_1) when

$$M = 0, \pm 1, \pm 2, \dots , \quad N = 0, \pm 1, \pm 2, \dots , \quad a = + | M | , \quad b = | K | . \quad (2.3a)$$

The wave function will be finite if the hypergeometric function becomes a polynomial:

$$A = -n, \quad n = 1, 2, 3, \dots ,$$

which provides us with the spectrum for energy:

$$\epsilon_N = \frac{1}{2} (N^2 - 1) , \quad N = a + b + 1 + 2n ; \quad (2.3b)$$

where the number N takes on the values $+1, +2, +3, \dots$. The energy level are degenerated as much as $g_N = N^2$. For instance, let $N = 3$ then $g_3 = 9$:

$$\begin{aligned} & a = 0 \quad b = 2 \quad M = 0 \quad K = \pm 2 \\ & a = 1 \quad b = 1 \quad M = \pm 1 \quad K = \pm 1 \quad n = 0 \\ & a = 2 \quad b = 0 \quad M = \pm 2 \quad K = 0 \\ & a = 0 \quad b = 0 \quad M = 0 \quad K = 0 \quad n = 1 . \end{aligned}$$

Now, let us show that the constructed wave functions


$$\Phi_{\epsilon MK} = C_{MK}^\epsilon e^{+iM\phi} e^{+iKz} \sin^{+|M|} \rho \cos^{+|K|} \rho F(A, B, C; \cos^2 \rho), \quad (2.4)$$

are continuous in space S_3 . The whole space $S_3 - G(\rho, \phi, z)$ let be divided into three parts: $G = (G_1 \cup G_2 \cup G_3)$:

$$G_1 = G(0 < \rho < \pi/2) , \quad G_2 = G(\rho = 0) , \quad G_3 = G(\rho = \pi/2) . \quad (2.5)$$

For the part G_1 which can be sketched graphically as


Fig 1 The domain G_1



here pairs $(A, A'), (B, B')$ and so on, and also $(1, 1', 1'', 1''')$ represent respectively only one point in S_3 . It is evident that integer values for M and K will lead us to have the same values of $\Phi_{\epsilon MK}$ in all identified points on the plane (ϕ, z) . So the wave functions are continuous on the G_1 -part $n_a = (n_0, n_i)$ of S_3 . The domain G_2 representing the closed curve

$$n_0 = \cos z, \quad n_1 = 0, \quad n_2 = 0, \quad n_3 = \sin z, \quad (2.6a)$$

can be described by the scheme




So that any function $f(\rho = 0, \phi, z)$ depending on the mute variable ϕ will be discontinuous in S_3 . Evidently, the wave function being specified on G_2

$$M \neq 0 \implies \Phi_{\epsilon MK} = 0 ;$$

$$M = 0 \implies \Phi_{\epsilon 0K} = e^{+iKz} F(A, B, C; 1) = \Phi(n) . \quad (2.6b)$$

is continuous about the curve G_2 . Analogously, for the domain G_3

$$n_0 = 0, \quad n_1 = \cos \phi, \quad n_2 = \sin \phi, \quad n_3 = 0 ; \quad (2.7a)$$



and

$$K \neq 0 \implies \Phi_{\epsilon MK} = 0$$

$$K = 0 \implies \Phi_{\epsilon M0} = e^{+iM\phi} F(A, B, C; 0) = \Phi(n) . \quad (2.7b)$$

Now let us consider the orthogonality condition for the wave function $\Phi_{\epsilon MK}(\rho, \phi, z)$. To this end, one should instead of ρ coordinate introduce another variable $x = \cos 2\rho$ and take into account defining relations for Yacobi polynomials [17]

$$P_n^{(a,b)}(x) = N_n^{(a,b)} F(-n, a+b+1+n, b+1; \frac{1+x}{2}) , \quad N_n^{(a,b)} = (-1) \frac{n(n+b)!}{b!} ,$$

then the functions $\Phi_{\epsilon MK}$ read as

$$\Phi_{\epsilon MK}(\rho, \phi, z) = \frac{1}{2\pi} e^{+iM\phi} e^{+iKz} \frac{C_{MK}^\epsilon}{N_n^{(a,b)}} \frac{1}{2^{a/2} 2^{b/2}} (1-x)^{a/2} (1+x)^{b/2} P_n^{(a,b)} . \quad (2.8a)$$

So we have relationship

$$\begin{aligned} \int_{S_3} \Phi_{\epsilon MK}^* \Phi_{\epsilon' M' K'} dV &= \delta_{MM'} \delta_{KK'} \frac{1}{2^{a+b}} \times , \\ &\times (C_{MK}^{\epsilon*} C_{M'K'}^{\epsilon'}) \frac{N_n^{(a,b)}}{N_{n'}^{(a',b')}} \frac{1}{4} \int_{-1}^{+1} (1-x)^a (1+x)^b P_n^{(a,b)} P_{n'}^{(a,b)} dx . \end{aligned} \quad (2.8b)$$

Taking in mind the orthogonality conditions for Yacobi polynomials [17]

$$\int_{-1}^{+1} (1-x)^a (1+x)^b P_n^{(a,b)} P_{n'}^{(a,b)} dx = \delta_{nn'} \frac{2^{a+b+1} (n+a)! (n+b)!}{n! (a+b+n)! (a+b+1+2n)} , \quad (2.8c)$$

from (2.8b) it follows

$$\int_{S_3} \Phi_{\epsilon MK}^* \Phi_{\epsilon' M' K'} dV = C \delta_{\epsilon\epsilon'} \delta_{MM'} \delta_{KK'} \quad (2.8d)$$

where C is defined by

$$C = \frac{|C_{MK}^\epsilon|^2}{|N_n^{(a,b)}|^2} \frac{(n+a)! (n+b)!}{2(a+b+1+2n)n!(a+b+n)} .$$

Demanding $C = 1$ we arrive at the normalized wave solutions – see (2.8d):

$$\Phi_{\epsilon MK} = \frac{e^{iM\phi}}{\sqrt{2\pi}} \frac{e^{iKz}}{\sqrt{2\pi}} A_n^{(a,b)} (1-x)^{a/2} (1+x)^{b/2} P_n^{(a,b)}(x) , \quad (2.9)$$

$$A_n^{(a,b)} = 2 \sqrt{\frac{n!(a+b+n)!(a+b+1+2n)}{(n+a)!(n+b)!2^{a+b+1}}} .$$

It is convenient to have the following table (restricting ourselves to $N = 1, 2, 3, 4$):

<u>$N=1$</u>	$b=0$	
$a=0$	$\sqrt{1} \ 1 \times$ $P_0^{(0,0)}(x)$	
<u>$N=2$</u>	$b=0$	$b=1$
$a=0$		$\sqrt{1 \ (1+x)^1} \times$ $P_0^{(0,1)}(x)$
$a=1$	$\sqrt{(1-x)^1 \ 1} \times$ $P_0^{(1,0)}(x)$	

<u>$N = 3$</u>	$b = 0$	$b = 1$	$b = 2$	
$a = 0$	$\sqrt{1-x} \times P_1^{(0,0)}(x)$		$\sqrt{1(1+x)^2} \times P_0^{(0,2)}(x)$	
$a = 1$		$\sqrt{(1-x)(1+x)} \times P_1^{(1,1)}(x)$		
$a = 2$	$\sqrt{(1-x)^2} \times P_0^{(2,0)}(x)$			
<u>$N = 4$</u>	$b = 0$	$b = 1$	$b = 2$	$b = 3$
$a = 0$		$\sqrt{1(1+x)} \times P_1^{(0,1)}(x)$		$\sqrt{1(1+x)^3} \times P_0^{(0,3)}(x)$
$a = 1$	$\sqrt{(1-x)^1} \times P_1^{(1,0)}(x)$		$\sqrt{(1-x)(1+x)^2} \times P_0^{(1,2)}(x)$	
$a = 2$		$\sqrt{(1-x)^2(1+x)} \times P_0^{(2,1)}(x)$		
$a = 3$	$\sqrt{(1-x)^3} \times P_0^{(3,0)}(x)$			

It should be given special attention to the following: the system of all $\{\Psi_{\epsilon MK}\}$ can be divided into two parts depending of N , even or odd. Orthogonality of the functions of opposite classes is achieved by integration over in (ϕ, z) -variables.

3. The case of elliptical space

Now let us turn to the elliptical case. It is a Riemmanian space of constant positive curvature as well, though different in its topological structure S_3 is 1-connected whereas $\tilde{S}_3$ is 2-connected. These space models have simple realizations as parametric manifolds of the unitary and orthogonal groups, $SU(2)$ and $S0(3)$:

$$B = \sigma^0 n_0 - i \sigma^k n_k, \quad \det B = +1, \quad (3.1a)$$

$$0(\vec{c}) = I + 2 \frac{\vec{c}^\times + (\vec{c}^\times)^2}{1 + \vec{c}^2}, \quad (\vec{c}^\times)_{kl} = -\epsilon_{klj} c_j. \quad (3.1b)$$

The matrix $O(\vec{c})$ represents a point in elliptic space. To every pair of different vector of infinite length $\vec{c}_\infty^\pm = \pm \infty \vec{c}_0$, $\vec{c}_0^2 = 1$ corresponds one the same matrix, point in elliptical space

$$0(\vec{c}^{\pm \infty}) = I + 2 (\vec{c}_0^\times)^2. \quad (3.1c)$$

The mapping $2 \implies 1$ from $SU(2)$ to $S0(3)$ $\{+n_a, -n_a\} \implies c^i = n_i/n_0$ will be used to establish a domain $\tilde{G}$ of cylindrical coordinates of elliptical model on the base of the known that for spherical one.

At searching the wave Schrödinger solutions in elliptical space one does not need to repeat all the calculation, instead it is sufficient to separate continuous solutions in space $\tilde{S}_3$ among these continuous in S_3 . So, we start with the relationships

$$c_1 = \frac{\tan \rho}{\cos z} \cos \phi, \quad c_2 = \frac{\tan \rho}{\cos z} \sin \phi, \quad c_3 = \tan z \quad (3.2a)$$

where the coordinates (ρ, ϕ, z) change in the limits:

$$\tilde{G}^{(+1)} = \{ \rho \in [0, \pi/2], \phi \in [-\pi, +\pi], z \in [-\pi/2, +\pi/2] \}. \quad (3.2b)$$

From the very beginning, one point should be emphasized: the domain $\tilde{G}$ according to (3.2b) is the most simple and evident variant, though not unique several other might be used as well. For instance, the following $\tilde{G}'$ is appropriate:

$$\tilde{G}^{(+1)} = \{ \rho \in [0, \pi/2], \phi \in [-\pi, +\pi], z \in [-\pi, -\pi/2] \oplus [+ \pi/2, +\pi] \}. \quad (3.2c)$$

Transition from elliptical curved model to a flat space model looks differently for cases $\tilde{G}^{(\delta=\pm 1)}$:

$$\underline{(\delta = +1)}$$

$$\rho \rightarrow 0, z \rightarrow \pm 0 : \quad c_1 = \rho \cos \phi, \quad c_2 = \rho \sin \phi, \quad c_3 = z ;$$

$$\underline{(\delta = -1)}$$


$$\rho \rightarrow 0, \quad z > 0, \quad z = \pi - Z \rightarrow \pi : \quad -c_1 = \rho \cos \phi, \quad -c_2 = \rho \sin \phi, \quad -c_3 = Z > 0 ;$$

$$\rho \rightarrow 0, \quad z < 0, \quad z = -\pi - Z \rightarrow \pi : \quad -c_1 = \rho \cos \phi, \quad -c_2 = \rho \sin \phi, \quad -c_3 = Z < 0 .$$

The variant with $\delta = +1$ is simpler and it will be used below³. One should find identification rules adjusted with topological structure of the group $SO(3.R)$. Again, the domain $\tilde{G}$ is to be divided into three parts:

$$\tilde{G}_1 = \tilde{G}(\rho \neq 0, \pi/2), \quad \tilde{G}_2 = \tilde{G}(\rho = 0), \quad \tilde{G}_3 = \tilde{G}(\rho = \pi/2) .$$

First, let us consider the domain $\tilde{G}_1$. The topological structure of the group $SO(3.R)$ leads us to the following identification scheme:




here (A, A') , (B, B') and so on, also $(1, 1', 1'')$ and $(2, 2', 2'')$ represent one the same respective points in $\tilde{S}_3$. Let us give some details.

³ However, these alternative variants should be taken in mind; in the following they will help to understand some peculiarity of tetrad formalism in elliptical space.

Consider the vicinity of the point $1'$; one can approach it by different directions

Fig 5 Vicinity of $1'$



The α -direction is described by the equation

$$\phi = (-\pi + \frac{\pi}{2} \tan \alpha) + z \tan(\pi - \alpha) . \quad (3.3a)$$

Near to $1'$ one has $z = (\pi/2 - \delta)$, δ is a very small positive number, then eq. (3.3a) reads

$$\phi = -\pi + \delta \tan \alpha , \cos \phi = -\cos(\delta \tan \alpha) , \sin \phi = -\sin(\delta \tan \alpha) ;$$

and therefore c^j in vicinity of $1'$ looks

$$c^1 = \tan \rho \frac{-\cos(\delta \tan \alpha)}{\sin \delta} , c^2 = \tan \rho \frac{-\sin(\delta \tan \alpha)}{\sin \delta} , c^3 = \frac{\cos \delta}{\sin \delta} . \quad (3.3b)$$

The limit of the above when $\delta \rightarrow 0$ ($0 \leq \alpha < \pi/2$) is one the same without any dependence of α :

$$\underline{1'} : \quad \vec{c} = \infty (-\tan \rho , 0 , 1) . \quad (3.3c)$$


Take notice that $\alpha \neq \pi/2$; Analogously for the vicinity of the point $1''$ one has

$$\underline{1''} : \quad \vec{c} = \infty (-\tan \rho , 0 , 1) . \quad (3.4)$$

And finally, for the point 1:

$$\phi = -\frac{\pi}{2} \tan g + z \tan(\pi - \alpha) ,$$

Fig 6 The vicinity of 1



here $z = (-\pi/2 + \delta)$ и $\phi = -\delta \tan \alpha$, and

$$c_1 = \tan \rho \frac{+\cos(\delta \tan \alpha)}{\sin \delta} , c_2 = \tan \rho \frac{-\sin(\delta \tan \alpha)}{\sin \delta} , c_3 = \frac{-\cos \delta}{\sin \delta} .$$

When δ changes to zero, one has

$$\underline{1} : \quad \vec{c} = -\infty (-\tan \rho , 0 , 1) . \quad (3.5)$$

Therefore, the $(1, 1', 1'')$ indeed represent one the same point in elliptical space. In the same manner, three points $(2, 2', 2'')$ correspond to


$$\underline{2} : \quad \vec{c} = +\infty (\tan \rho , 0 , 1) ,$$

$$\begin{aligned}\underline{2'}: \quad \vec{c} &= +\infty (\tan \rho, 0, 1), \\ \underline{2'':} \quad \vec{c} &= -\infty (\tan \rho, 0, 1).\end{aligned}\tag{3.6}$$

For instance. let us detail four other points:


$$\begin{aligned}\underline{A}: \quad \vec{c} &= +\infty (0, \tan \rho, 1), \quad \underline{A'}: \quad \vec{c} = (0, \tan \rho, 1), \\ \underline{B}: \quad \vec{c} &= +\infty (0, -\tan \rho, 1), \quad \underline{B'}: \quad \vec{c} = -\infty (0, -\tan \rho, 1).\end{aligned}\tag{3.7}$$

Identification on the boundaries $(1'CD2')$ and $(1''C'D'2'')$ is shown in the Fig 4; here only finite vector $\vec{c}$ are presented. Consideration of the boundary of $\tilde{G}_1$ is finished. Transition from $SU(2)$ to $SO(3.R)$ group spaces can be characterized by the scheme



Now we can examine two remaining domains. In the $\tilde{G}_2$ for right point P we have

$$\begin{aligned}\tilde{G}_2 \quad (\phi - \text{любое}), \\ \rho = \pi/2 \tan \alpha + z \tan \alpha \Rightarrow z = -\pi/2 + \delta, \rho = \delta \tan \alpha,\end{aligned}$$



Therefore,

$$c_1 = \frac{\tan(\delta \tan \alpha)}{\sin(\delta \tan \alpha)} \frac{\cos \phi}{\sin \delta}, \quad c_2 = \frac{\tan(\delta \tan \alpha)}{\sin(\delta \tan \alpha)} \frac{\sin \phi}{\sin \delta}, \quad c_3 = -\frac{\cos \delta}{\sin \delta},$$

in the limit $\delta \rightarrow 0$ ($\alpha \neq 0$) it follows

$$\underline{\tilde{G}_2}: \quad P, \quad \vec{c} = +\infty (0, 0, -1); \tag{3.8a}$$


In the same manner, for the left point P' we have

$$\underline{G_2}: \quad P', \quad \vec{c} = -\infty (0, 0, -1). \tag{3.8b}$$

So, all the domain $\tilde{G}_2$ determines the point of elliptical space in accordance with the formula

$$\underline{G_2}: \quad \vec{c} = -\infty (0, 0, \tan z) \tag{3.8c}$$

and ϕ is a mute variable. Mow consider third domain $\tilde{G}_3$:



Near the point B we have

$$\rho = -z \tan \alpha + \left(\frac{\pi}{2} - \frac{\pi}{2} \tan \alpha\right) \implies z = \left(-\frac{\pi}{2} + \delta\right), \quad \rho = \left(\frac{\pi}{2} - \delta \tan \alpha\right);$$

and

$$c_1 = \frac{\cos(\delta \tan \alpha)}{\sin(\delta \tan \alpha)} \frac{\cos \phi}{\sin \delta}, \quad c_2 = \frac{\cos(\delta \tan \alpha)}{\sin(\delta \tan \alpha)} \frac{\sin \phi}{\sin \delta}, \quad c_3 = -\frac{\cos \delta}{\sin \delta}.$$

From where, in the limit $\delta \rightarrow 0, \alpha \neq \pi/2$, it follows

$$\underline{B}: \quad \vec{c} = \infty (\cos \phi, \sin \phi, 0). \quad (3.9a)$$

In the same manner, for the point B' we have

$$\underline{B'}: \quad \vec{c} = \infty (\cos \phi, \sin \phi, +0). \quad (3.9b)$$

All the domain $\tilde{G}_3$ parameterizes the point in $\tilde{S}_3$ in accordance with the relation

$$\underline{B'}: \quad \vec{c} = \frac{\infty}{\cos z} (\cos \phi, \sin \phi, +\frac{\sin z}{\infty}) \sim \infty (\cos \phi, \sin \phi, 0), \quad (3.9c)$$

and z is a mute variable.

Now we are ready to establish which of the wave functions $\Phi_{\epsilon MK}$ founded in §2 are continuous in elliptical space and which are not. From the equation (see Fig 4)

$$f(1) = f(1') = f(1'')$$

one can reduce

$$e^{-iK(\pi/2)} = e^{-iM\pi} e^{+iK(\pi/2)} = e^{+iM\pi} e^{+iK(\pi/2)},$$

from where it follows

$$e^{i2M\pi} = 1, \quad e^{i(K-M)\pi} = 1, \quad e^{i(K+M)\pi} = 1;$$

that is $(K - M)$ and $(K + M)$ must be integer and even. Analogous relations for the points $(2, 2', 2'')$ results in the same.

Therefore, the functions $\phi_{\epsilon MK}(\rho, \phi, z)$ founded in §2 will be single-valued and continuous in the $\tilde{G}_1$ if M and K are both even or both odd. Correspondingly, the main quantum number N takes on the values $N = 1, 3, 5, \dots$

In the domain $\tilde{G}_2$ the function $\Phi_{\epsilon KM}(\rho, \phi, z)$ given by

$$G_2 : \quad \phi_{\epsilon MK} = \begin{cases} 0, & M \neq 0; \\ e^{iKz} F(A, B, C; 1), & M = 0 \end{cases}, \quad (3.10)$$

is a continuous function on $\tilde{S}_3$. In the $\tilde{G}_3$ we have behavior correct in the sense of continuity as well

$$\tilde{G}_3 = \begin{cases} 0, & K \neq 0, \\ e^{iMz} F(A, B, C; 0), & K = 0 \end{cases}. \quad (3.11)$$

Thus, the functions $\Phi_{\epsilon MK}(\rho, \phi, z)$, $N = 1, 3, 5, \dots$ provide us with the system of single-valued and continuous in the elliptical space $SO(3.R)$ – one formal change must be done:

$$\frac{1}{\sqrt{2\pi}} e^{iKz} \implies \frac{1}{\sqrt{\pi}} e^{iKz},$$

$$\int_{SO(3)} \Phi_{\epsilon MK}^* \Phi_{\epsilon' M' K'} dV = \delta_{MM'} \delta_{KK'} \delta_{NN'}, \quad N = 1, 3, 5, \dots \quad (3.12)$$

At the same time, the remaining functions

$$\Phi_{\epsilon MK}(\rho, \phi, z), \quad N = 2, 4, \dots \quad (3.13)$$

are discontinuous in elliptical space $SO(3.R)$ and they do not consist of a basis in the space of continuous function on the $SO(3)$ though obey the same orthogonality conditions as (3.12).

4. Euler variables as coordinates on the group $SU(2)$.

Now we are to refer the above Schrödinger's wave solutions $\Phi_{\epsilon MK}(\rho, \phi, z)$ to the known Wigner's D -functions [17].

$$\begin{aligned} \Phi_{\epsilon MK} &= \frac{e^{iM\phi}}{\sqrt{2\pi}} \frac{e^{iK\phi}}{\sqrt{2\pi}} \left[A_n^{(a,b)} (1-x)^{a/2} (1+x)^{b/2} P_n^{(a,b)}(x) \right] = \\ &= \frac{e^{iM\phi}}{\sqrt{2\pi}} \frac{e^{iK\phi}}{\sqrt{2\pi}} d_{m\sigma}^j(\beta) = D_{m\sigma}^j(\alpha, \beta, \gamma); \end{aligned} \quad (4.1a)$$

where

$$\begin{aligned} N &= (2j+1), \quad \epsilon_N = \frac{1}{2}(N^2 - 1) = 2j(j+1), \\ m &= -\frac{1}{2}(M+K), \quad \sigma = -\frac{1}{2}(M-K), \end{aligned} \quad (4.1b)$$

and coordinates (ρ, ϕ, z) are connected with Euler variables (α, β, γ) as follows:

$$\alpha = \phi + z, \quad \beta = 2\rho, \quad \gamma = \phi - z. \quad (4.1c)$$

This transition can be traced also in differential equations

$$\begin{aligned} &[\sin^{-1} \rho \cos^{-1} \rho \partial_\rho \sin \rho \cos \rho \partial_\rho + \sin^{-2} \rho \partial_\phi^2 + \cos^{-2} \rho \partial_z^2 + 2\epsilon] \Phi = 0, \\ &-i\partial_\phi \Phi = M \Phi, \quad -i\partial_z \Phi = K \Phi \end{aligned} \quad (4.2a)$$

which in (α, β, γ) variables become

$$[-\sin^{-1} \beta \partial_\beta \sin \beta \partial_\beta + \sin^{-2} \beta (-\partial_\alpha^2 + 2 \cos \beta \partial_\alpha \partial_\gamma - \partial_\gamma^2)] \Phi = (\epsilon/2) \Phi,$$

$$i\partial_\alpha \Phi = -\frac{1}{2}(M+K) \Phi, \quad i\partial_\gamma \Phi = -\frac{1}{2}(M-K) \Phi. \quad (4.2b)$$

and coincide with determining relations for D -functions: $\Phi_{\epsilon MK}(\rho, \phi, z) = D_{m\sigma}^j(\alpha, \beta, \gamma)$.


All specifics of the Euler angles as parameters on the unitary group can be found with absolute fullness from the previous analysis. First, these variables are non-orthogonal coordinates in S_3 :

$$dl^2 = \frac{1}{4} [d\beta^2 + d\alpha^2 - 2 \cos \beta d\alpha d\gamma + d\gamma^2]. \quad (4.2c)$$

Again, the whole domain $G(\alpha, \beta, \gamma)$ should be divided into three parts $G = (G_1 \cup G_2 \cup G_3)$:

$$G_1 = G(\beta \neq 0, \pi), \quad G_2 = G(\beta = 0), \quad G_3 = G(\beta = \pi). \quad (4.3)$$


The domain $G_1(\alpha, \beta)$ is characterized by the scheme



The closed curve associated with the domain $G_2(\alpha, \gamma)$

$$n_0 = \cos \frac{\alpha - \gamma}{2}, \quad n_1 = 0, \quad n_2 = 0, \quad n_3 = \sin \frac{\alpha - \gamma}{2}; \quad (4.4a)$$

from where it follows that (α, β) and (α', β') represent the same point in the curve if $(\alpha - \gamma) = (\alpha' - \beta')$. In accordance with this the domain $G_2(\alpha, \gamma)$ is characterized by the scheme




here the full segment $(\alpha, \beta)_z$ given by equation $\alpha = \gamma + 2z$ corresponds to the single point $n = (\cos z, 0, 0, \sin z)$. Wigner functions in the $G_2(\alpha, \gamma)$

$$D_{m\sigma}^j(\alpha, 0, \gamma) = \begin{cases} 0, & (m + \sigma) \neq 0, \\ F(A, B, C; 1), & m = 0, \sigma = 0, \\ e^{im(\alpha - \gamma)} F(A, B, C; 1), & (m + \sigma) = 0 \end{cases} \quad (4.4b)$$

are continuous in the curve. Consideration of the domain G_3 will be the same:

$$n_0 = 0, \quad n_1 = \cos \frac{\alpha + \gamma}{2}, \quad n_2 = \sin \frac{\alpha + \gamma}{2}, \quad n_3 = 0; \quad (4.5a)$$

Fig 12 $G_3(\alpha, \gamma)$



and Wigner function


$$D_{m\sigma}^j(\alpha, \pi, \gamma) = \begin{cases} 0, & (m - \sigma) \neq 0, \\ F(A, B, C; 0), & m = 0, \sigma = 0, \\ e^{-im(\alpha + \gamma)} F(A, B, C; 0), & (m - \sigma) = 0; \end{cases} \quad (4.5b)$$

depending on the argument $(\alpha + \beta)$ are continuous in the curve.

Now let us reformulate the orthogonality conditions (2.8):

$$\int_{S_3} \Phi_{\epsilon MK}^* \Phi_{\epsilon' M' K'} dV = \delta_{\epsilon \epsilon'} \delta_{MM'} \delta_{KK'}, \quad dV = \sin \rho \cos \rho d\rho d\phi dz,$$

Fig 13 $G(\phi, z)$




which after transition to Euler variable will look as follows:

$$\int_{SU(2)} D_{m\sigma}^{j*}(\alpha, \beta, \gamma) D_{m'\sigma'}^{j'}(\alpha, \beta, \gamma) d\omega = \delta_{mm'} \delta_{\sigma\sigma'} \delta_{jj'},$$

$$d\omega = \frac{1}{8} \sin \beta d\alpha d\beta d\gamma; \quad 0 \leq \beta \leq \pi,$$

Fig 14 $G(\alpha, \gamma)$



Indeed, taking the following order in integrating

$$\begin{aligned} & \int_{G(\alpha, \gamma)} d\alpha d\gamma F(\alpha, \gamma) = \\ & = \int_{-2\pi}^0 d\gamma \int_{\alpha=-(\gamma+2\pi)}^{\alpha=+(\gamma+2\pi)} d\alpha F(\alpha, \gamma) + \int_0^{+2\pi} d\gamma \int_{\alpha=-(-\gamma+2\pi)}^{\alpha=+(-\gamma+2\pi)} d\alpha F(\alpha, \gamma) . \end{aligned} \quad (4.6)$$

it is the matter of simple calculation to obtain the identity

$$\frac{1}{8\pi^2} \int_{G(\alpha, \gamma)} d\alpha d\gamma [e^{i(m-m')\alpha} e^{i(\sigma-\sigma')\gamma}] = \delta_{mm'} \delta_{\sigma\sigma'} , \quad (4.7)$$

One can easily see that the same symbol $(\delta_{mm'} \delta_{\sigma\sigma'})$ might be achieved in integrating over two other domains, $\Delta(\alpha, \gamma)$ or $\Delta'(\alpha, \gamma)$ determined in Fig 15 and Fig 16 below.

Fig 15 $\Delta(\alpha, \gamma)$





Fig 16 $\Delta'(\alpha, \gamma)$



This fact is not occasional. Indeed, now we will show that a more general statement is true: *if $\Phi(\alpha, \beta, \gamma) = \Phi(n)$ is any single-valued and continuous function on the sphere S_3 , then identity holds (or $\Delta \rightarrow \Delta'$)*

$$\int_{R(\beta) \otimes G(\alpha, \gamma)} d\omega \Phi(n) = \int_{R(\beta) \otimes \Delta(\alpha, \gamma)} d\omega \Phi(n) . \quad (4.8)$$

Indeed, any that function $\Phi(n)$, being expressed in coordinates (ρ, ϕ, z) , gives a function 2π -periodical in ϕ и z :

$$\Phi(\rho, \phi', z') = \Phi(\rho, \phi'', z'') \iff (\phi'' - \phi') = 2\pi\nu , \quad (z' - z'') = 2\pi\mu . \quad (4.9a)$$

After translating to variables (α, γ) one has

$$\frac{1}{2} (\alpha - \gamma) = \phi , \quad \frac{1}{2} (\alpha + \gamma) = z ,$$


$$F(\rho, \alpha, \gamma) = \Phi(\rho, \frac{\alpha + \gamma}{2}, \frac{\alpha - \gamma}{2}) .$$

Therefore, any function on S_3 , single-valued and continuous, must obey the following periodicity-condition:

$$\begin{aligned} F(\rho, \alpha', \gamma') &= F(\rho, \alpha'', \gamma'') , \\ (\alpha'' - \alpha') &= 2\pi(\nu + \mu) , \quad (\gamma'' - \gamma') = 2\pi(\nu - \mu) . \end{aligned} \quad (4.9b)$$

Now let us compare two domains, $G(\alpha, \gamma)$ and $\Delta(\alpha, \gamma)$:

Fig 17



It is easily seen that the parts A, B, C, D and A', B', C', D' can be moved to each other by respective shifts:


$$\begin{aligned} A &\implies A' (\alpha' = \alpha - 2\pi, \gamma' = \gamma + 2\pi) , \\ B &\implies B' (\alpha' = \alpha - 2\pi, \gamma' = \gamma - 2\pi) , \\ C &\implies C' (\alpha' = \alpha + 2\pi, \gamma' = \gamma + 2\pi) , \\ D &\implies D' (\alpha' = \alpha + 2\pi, \gamma' = \gamma - 2\pi) . \end{aligned}$$

From where, taking in mind

$$G = [G_0 + (A + B + C + D)] , \quad \Delta = [G + (A' + B' + C' + D')]$$

and the above 2π -periodicity of $F(\rho, \alpha, \gamma)$ in variables α and γ , one arrives at (4.8). Consideration of the domain Δ' is the same.

Thus, to cover the group $SU(2)$, instead of $[R(\rho) \otimes G(\alpha, \gamma)]$, one may use other variants, $[R(\rho) \otimes \Delta(\alpha, \gamma)]$ or $[R(\rho) \otimes \Delta'(\alpha, \gamma)]$. However, one should remember on changing the identification rule for the boundary points. For instance, tracing the all identical points in the replacement $G \rightarrow \Delta$ we get a new scheme:




It formally coincides with that for the cylindrical variables (ϕ, z) in the case of orthogonal group $SO(3.R)$ group.

5. Euler angles, coordinates on the group $S0(3.R)$.

Transition to the Euler variables in the theory of orthogonal group can be done as before. Below let us written down only the main graphical schemes.


Instead of Fig. 9 there is

Fig 19 $G_1(\alpha, \gamma)$




Instead of Fig 11 there is

Fig 20 $G_2(\alpha, \gamma)$ $\vec{c} = (0, 0, \tan z)$




Instead of Fig 12 there is

Fig 21 $\vec{c} = \infty (\cos \phi, \sin, 0)$




Instead of Fig 17

Fig 22 $G(\alpha, \gamma) \rightarrow \Delta(\alpha, \gamma)$



Instead of Fig 18 there is

Fig 23 $\Delta_1(\alpha, \gamma)$



PART II. DIRAC EQUATION ON THE SPHERE S_3

6. Dirac equation in cylindrical coordinates, separating the variables

Let us turn to the Dirac equation in cylindrical coordinates and tetrad on the sphere S_3 :

$$\begin{aligned}
 n_1 &= \sin \rho \cos \phi, \quad n_2 = \sin \rho \sin \phi, \\
 n_3 &= \cos \rho \sin z, \quad n_4 = \cos \rho \cos z; \\
 dl^2(y) &= d\rho^2 + \sin^2 \rho d\phi^2 + \cos^2 \rho dz^2, \\
 e_{(a)}^\alpha(y) &= \begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \sin^{-1} \rho & 0 \\ 0 & 0 & 0 & \cos^{-1} \rho \end{vmatrix}; \quad (6.1)
 \end{aligned}$$

where (ρ, ϕ, z) change in the domain

$$G = \{ \rho \in [0, +\pi/2], \quad \phi \in [-\pi, +\pi], \quad z \in [-\pi, +\pi] \}.$$

As in scalar case, here it is convenient to divide $G(\rho, \phi, z)$ into three parts: $G_1(\rho \neq 0, \pi/2)$, $G_2(\rho = 0)$, $G_3(\rho = \pi/2)$. The Dirac equation in orthogonal coordinates and tetrad can be written in the form

$$[i \gamma^a (e_{(a)}^\alpha \frac{\partial}{\partial x^\alpha} + \frac{1}{2} (\frac{1}{\sqrt{-g}} \frac{\partial}{\partial x^\alpha} \sqrt{-g} e_{(a)}^\alpha)) - M] \Psi(x) = 0. \quad (6.2a)$$

At the given tetrad (6.1) equation (6.2a) will read

$$\left[\gamma^0 \frac{\partial}{\partial t} + \gamma^1 \left(\frac{\partial}{\partial \rho} + \frac{\cos 2\rho}{\sin 2\rho} \right) + \gamma^2 \frac{1}{\sin \rho} \frac{\partial}{\partial \phi} + \gamma^3 \frac{1}{\cos \rho} \frac{\partial}{\partial z} + iM \right] \Psi = 0. \quad (6.2b)$$

Let us construct solutions, the eigenfunctions of $i\partial_t$, $i\partial_\phi$, $i\partial_z$:

$$\Psi(t; \rho, \phi, z) = e^{-i\epsilon t} e^{im\phi} e^{ikz} \begin{vmatrix} f_1(\rho) \\ f_2(\rho) \\ f_3(\rho) \\ f_4(\rho) \end{vmatrix}. \quad (6.3)$$

With the use of explicit form of the Dirac matrices in the spinor basis from (6.2b) it follows equations for f_i :

$$\begin{aligned} -i\epsilon f_3 - \left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_4 - \frac{m}{\sin \rho} f_4 - \frac{ik}{\cos \rho} f_3 + iM f_1 &= 0, \\ -i\epsilon f_4 - \left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_3 + \frac{m}{\sin \rho} f_3 + \frac{ik}{\cos \rho} f_4 + iM f_2 &= 0, \\ -i\epsilon f_1 + \left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_2 + \frac{m}{\sin \rho} f_2 + \frac{ik}{\cos \rho} f_1 + iM f_3 &= 0, \\ -i\epsilon f_2 + \left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_1 - \frac{m}{\sin \rho} f_1 - \frac{ik}{\cos \rho} f_2 + iM f_4 &= 0. \end{aligned} \quad (6.4)$$

To simplify the system (6.4) one other operator should be diagonalized. In flat space, as that can be taken the helicity operator

$$(\vec{\Sigma} \vec{P}) \Psi_{cart} = \lambda \Psi_{cart}, \quad \vec{P} = -i\vec{\nabla}.$$

Because the cartesian and cylindrical bases are related by spinor gauge transformation over fermion wave functions:

$$\Psi^{cyl} = S \Psi^{cart}, \quad S = \begin{vmatrix} B & 0 \\ 0 & B \end{vmatrix}, \quad B = \begin{vmatrix} e^{+i\phi/2} & 0 \\ 0 & e^{-i\phi/2} \end{vmatrix}, \quad (6.5a)$$

for the helicity operator in cylindrical representation

$$\begin{aligned} B \vec{\Sigma} \vec{P} B^{-1} &= \begin{vmatrix} P_3 & e^{+i\phi/2} (P_1 - iP_2) e^{-i\phi/2} \\ e^{+i\phi/2} (P_1 + iP_2) e^{-i\phi/2} & -P_3 \end{vmatrix}, \\ (P_1 \pm iP_2) &= -ie^{\pm i\phi/2} \left(\frac{\partial}{\partial \rho} \pm \frac{i}{\rho} \frac{\partial}{\partial \phi} \right), \\ e^{\mp i\phi/2} (P_1 \pm iP_2) e^{\pm i\phi/2} &= \left[-i \left(\frac{\partial}{\partial \rho} + \frac{1}{2\rho} \right) \pm \frac{1}{\rho} \frac{\partial}{\partial \phi} \right], \end{aligned} \quad (6.5b)$$

one produces the expression

$$\Lambda^0 = \gamma^2 \gamma^3 \left(\frac{\partial}{\partial \rho} + \frac{1}{2\rho} \right) + \gamma^3 \gamma^1 \frac{1}{\rho} \frac{\partial}{\partial \phi} + \gamma^1 \gamma^2 \frac{\partial}{\partial z}. \quad (6.6)$$

Now, taking in mind eq. (6.6) and the Dirac equation (6.2b) in space S_3 , one may guess the form of a generalized helicity operator Λ in the curved space:

$$\Lambda = \gamma^2 \gamma^3 \left(\frac{\partial}{\partial \rho} + \frac{\cos 2\rho}{\sin 2\rho} \right) + \gamma^3 \gamma^1 \frac{1}{\sin \rho} \frac{\partial}{\partial \phi} + \gamma^1 \gamma^2 \frac{1}{\cos \rho} \frac{\partial}{\partial z}. \quad (6.7)$$

Equation on the proper values $\Lambda \Psi = \lambda \Psi$ leads us to

$$\begin{aligned}
-i\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_2 - \frac{im}{\sin \rho} f_2 + \frac{k}{\cos \rho} f_1 &= \lambda f_1 , \\
-i\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_1 + \frac{im}{\sin \rho} f_1 - \frac{k}{\cos \rho} f_2 &= \lambda f_2 , \\
-i\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_4 - \frac{im}{\sin \rho} f_4 + \frac{k}{\cos \rho} f_3 &= \lambda f_3 , \\
-i\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho}\right) f_3 + \frac{im}{\sin \rho} f_3 - \frac{k}{\cos \rho} f_4 &= \lambda f_4 .
\end{aligned} \tag{6.8}$$

Considering eqs. (6.8) and (6.4) together, one produces a linear algebraic system

$$\begin{aligned}
-i\epsilon f_3 - i\lambda f_3 + iM f_1 &= 0 , \\
-i\epsilon f_4 - i\lambda f_4 + iM f_2 &= 0 , \\
-i\epsilon f_1 + i\lambda f_1 + iM f_3 &= 0 , \\
-i\epsilon f_2 + i\lambda f_2 + iM f_4 &= 0 .
\end{aligned} \tag{6.9a}$$

From (6.9a) it follows

$$\lambda = \pm \sqrt{\epsilon^2 - M^2} , \quad f_3 = \frac{\epsilon - \lambda}{M} f_1 , \quad f_4 = \frac{\epsilon - \lambda}{M} f_2 ,$$

$$\Psi(t; \rho, \phi, z) = e^{-iet} e^{im\phi} e^{ikz} \begin{vmatrix} f_1(\rho) \\ f_2(\rho) \\ f_3(\rho) \\ f_4(\rho) \end{vmatrix} . \tag{6.9b}$$

Taking into account (6.9b), four equations in (6.4) give a simpler system for f_1, f_2 :

$$\begin{aligned}
\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} + \frac{m}{\sin \rho}\right) f_2 - i\left(\lambda - \frac{k}{\cos \rho}\right) f_1 &= 0 , \\
\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} - \frac{m}{\sin \rho}\right) f_1 - i\left(\lambda + \frac{k}{\cos \rho}\right) f_2 &= 0 .
\end{aligned} \tag{6.10}$$

By means of separating a special factor

$$f_1(\rho) = \frac{1}{\sqrt{\cos \rho \sin \rho}} \varphi_1(\rho) , \quad f_2(\rho) = \frac{1}{\sqrt{\cos \rho \sin \rho}} \varphi_2(\rho) ,$$

one can exclude the term $(\cos 2\rho / \sin 2\rho)$ from eqs. (6.10):

$$\begin{aligned}
\left(\frac{d}{d\rho} + \frac{m}{\sin \rho}\right) \varphi_2 - i\left(\lambda - \frac{k}{\cos \rho}\right) \varphi_1 &= 0 , \\
\left(\frac{d}{d\rho} - \frac{m}{\sin \rho}\right) \varphi_1 - i\left(\lambda + \frac{k}{\cos \rho}\right) \varphi_2 &= 0 .
\end{aligned} \tag{6.11}$$

7. Some additional transforming

Summing and subtracting equations in (6.11), one gets

$$\begin{aligned} \frac{d}{d\rho}(\varphi_1 + \varphi_2) - \frac{m}{\sin \rho}(\varphi_1 - \varphi_2) - i \lambda (\varphi_1 + \varphi_2) + \frac{k}{\cos \rho}(\varphi_1 - \varphi_2) &= 0 , \\ \frac{d}{d\rho}(\varphi_1 - \varphi_2) - \frac{m}{\sin \rho}(\varphi_1 + \varphi_2) + i \lambda (\varphi_1 - \varphi_2) - \frac{k}{\cos \rho}(\varphi_1 + \varphi_2) &= 0 . \end{aligned} \quad (7.1)$$

Now, instead of functions (φ_1, φ_2) it will be helpful to define new ones (F_1, F_2) ⁴:

$$\varphi_1 = \cos \frac{\rho}{2} F_1 - i \sin \frac{\rho}{2} F_2 , \quad \varphi_2 = -i \sin \frac{\rho}{2} F_1 + \cos \frac{\rho}{2} F_2 , \quad (7.2a)$$

$$(\varphi_1 + \varphi_2) = e^{-i\rho/2}(F_1 + F_2) , \quad (\varphi_1 - \varphi_2) = e^{+i\rho/2}(F_1 - F_2) . \quad (7.2b)$$

With (7.2b), eqs. (7.1) give

$$\frac{d}{d\rho}(F_1 + F_2) - \frac{i}{2}(F_1 + F_2) - \frac{m}{\sin \rho}e^{+i\rho}(F_1 - F_2) - i \lambda (F_1 + F_2) + \frac{ik}{\cos \rho}e^{+i\rho}(F_1 - F_2) = 0 ,$$

$$\frac{d}{d\rho}(F_1 - F_2) + \frac{i}{2}(F_1 - F_2) - \frac{m}{\sin \rho}e^{-i\rho}(F_1 + F_2) + i \lambda (F_1 - F_2) - \frac{ik}{\cos \rho}e^{-i\rho}(F_1 + F_2) = 0 .$$

Summing and subtracting them we will arrive at

$$\begin{aligned} \left(\frac{d}{d\rho} - m \frac{\cos \rho}{\sin \rho} - k \frac{\sin \rho}{\cos \rho} \right) F_1 &= i \left(\lambda + \frac{1}{2} - m + k \right) F_2 , \\ \left(\frac{d}{d\rho} + m \frac{\cos \rho}{\sin \rho} + k \frac{\sin \rho}{\cos \rho} \right) F_2 &= i \left(\lambda + \frac{1}{2} + m - k \right) F_1 . \end{aligned} \quad (7.3)$$

From (7.3), for $F_1(\rho)$ can be produced the second order equations: (take notice of the symmetry $m \Rightarrow -m, k \Rightarrow -k$):

$$\left[\frac{d^2}{d\rho^2} - \frac{m(m-1)}{\sin^2 \rho} - \frac{k(k+1)}{\cos^2 \rho} + \left(\lambda + \frac{1}{2} \right)^2 \right] F_1(\rho) = 0 , \quad (7.4a)$$

$$\left[\frac{d^2}{d\rho^2} - \frac{m(m+1)}{\sin^2 \rho} - \frac{k(k-1)}{\cos^2 \rho} + \left(\lambda + \frac{1}{2} \right)^2 \right] F_2(\rho) = 0 . \quad (7.4b)$$

One can again turn to the old functions

$$\begin{aligned} F_1(\rho) &= \sqrt{\sin \rho \cos \rho} G_1(\rho) , \\ F_2(\rho) &= \sqrt{\sin \rho \cos \rho} G_2(\rho) , \end{aligned} \quad (7.5)$$

then, taking in mind identity

$$\frac{d^2}{d\rho^2} \sqrt{\sin 2\rho} G = \sqrt{\sin 2\rho} \left[\frac{d^2}{d\rho^2} + 2 \frac{\cos 2\rho}{\sin 2\rho} \frac{d}{d\rho} - 1 - \frac{1}{\sin^2 2\rho} \right] G .$$

⁴Author is grateful to V.S. Otchik for pointing out this additional transformation over functions φ_1, φ_2 .

one produces

$$\left[\frac{d^2}{d\rho^2} + 2 \frac{\cos 2\rho}{\sin 2\rho} \frac{d}{d\rho} - 1 - \frac{1}{\sin^2 2\rho} - \frac{m(m-1)}{\sin^2 \rho} - \frac{k(k+1)}{\cos^2 \rho} + (\lambda + \frac{1}{2})^2 \right] G_1 = 0 , \quad (7.6a)$$

$$\left[\frac{d^2}{d\rho^2} + 2 \frac{\cos 2\rho}{\sin 2\rho} \frac{d}{d\rho} - 1 - \frac{1}{\sin^2 2\rho} - \frac{m(m+1)}{\sin^2 \rho} - \frac{k(k-1)}{\cos^2 \rho} + (\lambda + \frac{1}{2})^2 \right] G_2 = 0 . \quad (7.6b)$$

Again one can note the symmetry $m \Rightarrow -m$, $k \Rightarrow -k$. These equation can be rewritten as follows:

$$\left[\frac{d^2}{d\rho^2} + 2 \frac{\cos 2\rho}{\sin 2\rho} \frac{d}{d\rho} - \frac{(m-1/2)^2}{\sin^2 \rho} - \frac{(k+1/2)^2}{\cos^2 \rho} + (\lambda + \frac{1}{2})^2 - 1 \right] G_1 = 0 , \quad (7.7a)$$

$$\left[\frac{d^2}{d\rho^2} + 2 \frac{\cos 2\rho}{\sin 2\rho} \frac{d}{d\rho} - \frac{(m+1/2)^2}{\sin^2 \rho} - \frac{(k-1/2)^2}{\cos^2 \rho} + (\lambda + \frac{1}{2})^2 - 1 \right] G_2 = 0 . \quad (7.7b)$$

The corresponding first order equations look as

$$\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} - m \frac{\cos \rho}{\sin \rho} - k \frac{\sin \rho}{\cos \rho} \right) G_1 = i \left(\lambda + \frac{1}{2} - m + k \right) G_2 , \quad (7.8a)$$

$$\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} + m \frac{\cos \rho}{\sin \rho} + k \frac{\sin \rho}{\cos \rho} \right) G_2 = i \left(\lambda + \frac{1}{2} + m - k \right) G_1 . \quad (7.8b)$$

The initial radial functions $f_1(\rho)$, $f_2(\rho)$ are related with these G_1, G_2 by the formulas:

$$f_1(\rho) = \cos \frac{\rho}{2} G_1 - i \sin \frac{\rho}{2} G_2 , \quad f_2(\rho) = -i \sin \frac{\rho}{2} G_1 + \cos \frac{\rho}{2} G_2 . \quad (7.9)$$

$$\lambda = \pm \sqrt{\epsilon^2 - M^2} , \quad f_3 = M^{-1}(\epsilon - \lambda) f_1 , \quad f_4 = M^{-1}(\epsilon - \lambda) f_2 ,$$

$$\Psi_{\epsilon, m, k, \lambda} = e^{-i\epsilon t} e^{im\phi} e^{ikz} \begin{vmatrix} f_1(\rho) \\ f_2(\rho) \\ f_3(\rho) \\ f_4(\rho) \end{vmatrix} . \quad (7.10)$$

In the following it suffices to trace only by two first components of the fermion wave function:

$$\Psi_{\epsilon, m, k, \lambda} = \frac{e^{im\phi} e^{ikz}}{\sqrt{2}} \begin{vmatrix} \sqrt{1 + \cos \rho} G_1 - i \sqrt{1 - \cos \rho} G_2 \\ -i \sqrt{1 - \cos \rho} G_1 + \sqrt{1 + \cos \rho} G_2 \end{vmatrix} . \quad (7.11)$$

When ρ approaches the values 0 and $\pi/2$ the wave function ψ reads respectively

$$\underline{\rho \rightarrow 0} \quad \Psi_{\epsilon, m, k, \lambda} \rightarrow e^{im\phi} e^{ikz} \begin{vmatrix} G_1(0) - \frac{i\rho}{2} G_2(0) \\ -\frac{i\rho}{2} G_1(0) + G_2(0) \end{vmatrix} , \quad (7.12a)$$

$$\underline{\rho \rightarrow \pi/2} \quad \Psi_{\epsilon, m, k, \lambda} \rightarrow e^{im\phi} e^{ikz} \begin{vmatrix} G_1(\pi/2) - i G_2(\pi/2) \\ -i G_1(\pi/2) + G_2(\pi/2) \end{vmatrix} . \quad (7.12b)$$

8. Fermion in a curved space, tetrad gauge symmetry and formulation of continuity criterion for Dirac wave functions in spherical space S_3

The requirement of continuity for a fermion wave function in any Riemannian space-time is much more complicated than in the flat space-time by Minkowski. The matter is that the form of generally covariant Dirac equation [...] itself involves the tetrad formalism. In other words, one may use any form of the Dirac equation in space-time with a given metrical tensor $g_{\alpha\beta}(x)$ from many different ones depending on a chosen tetrad, all such tetrad-based fermion wave functions $\Psi_{tetrad}(x)$ are related to each other by local gauge transformations from the group $SL(2.C)$:

$$\Psi_{tetrad'}(x) = S(x) \Psi_{tetrad}(x), \quad S(x) \in SL(2.C) .$$

As a rule, these (4×4) gauge matrices $S(x)$ are themselves singular. So the explicit embodiment of the general continuity criterion for fermion wave functions depend on the occasional choice of the tetrad.

For more clarity, one may turn to the familiar case of flat space. It is tacitely accepted that in ordinary representation of the Dirac equation all fermion wave function must be single-valued and continuous. However, if one wishes to employ any other tetrad ⁵, for instance let it be the cylindrical tetrad (see (6.5a))

In the space E_3

$$\Psi^{cyl} = S \Psi^{cart}, \quad S = \begin{vmatrix} B & 0 \\ 0 & B \end{vmatrix}, \quad B = \begin{vmatrix} e^{+i\phi/2} & 0 \\ 0 & e^{-i\phi/2} \end{vmatrix}, \quad (8.1a)$$

because of the singular behavior of the gauge matrix $B(\phi)$ at the axis z the wave function Ψ^{cyl} will become discontinuous at the axis z :

$$\Psi_{cyl} = e^{-i\epsilon t} e^{ikz} \begin{vmatrix} e^{im\phi} f_1(\rho) \\ e^{im\phi} f_2(\rho) \\ e^{im\phi} f_3(\rho) \\ e^{im\phi} f_4(\rho) \end{vmatrix}, \quad \Psi_{cart} = e^{-i\epsilon t} e^{ikz} \begin{vmatrix} e^{im\phi} e^{-i\phi/2} f_1(\rho) \\ e^{im\phi} e^{+i\phi/2} f_2(\rho) \\ e^{im\phi} e^{-i\phi/2} f_3(\rho) \\ e^{im\phi} e^{+i\phi/2} f_4(\rho) \end{vmatrix}. \quad (8.1b)$$

From (8.1b) one may see that to the single-valued functions in Cartesian basin correspond the non-single-valued ones in the cylindrical basis. In eq. (8.1b) the quantum number m must be half-integer, one should note the gauge invariant character of the proper value problem:

$$\begin{aligned} cart : \quad \hat{J}_3 &= \hat{l}_3 + \hat{S}_3; \quad \hat{J}_3 \Psi_{Cart} = m \Psi_{art}, \\ cyl : \quad \hat{l}_3 \Psi_{cyl} &= m \Psi_{cyl}. \end{aligned} \quad (8.2)$$

In the curved space S_3 all the situation is principally the same though more complicated. There exists an analog of the Cartesian basis, here its role plays a conformally-flat basis (coordinates (x^1, x^2, x^3) and the tetrad $e_{(a)}^\alpha(x)$)

$$\begin{aligned} n_0^2 + n_1^2 + n_2^2 + n_3^2 &= 1, \\ x^1 &= \frac{n_1}{1 + n_4}, \quad x^2 = \frac{n_2}{1 + n_4}, \quad x^3 = \frac{n_3}{1 + n_4}, \end{aligned}$$

⁵A more detailed treatment see in [22,23].

$$dl^2(x) = \frac{1}{f^2} [(dx^1)^2 + (dx^2)^2 + (dx^3)^2] , \quad f = \frac{1+x^2}{2} ,$$

$$e_{(a)}^\alpha(x) = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & 1/f & 0 & 0 \\ 0 & 0 & 1/f & 0 \\ 0 & 0 & 0 & 1/f \end{vmatrix} . \quad (8.3a)$$

Spinor gauge transformation relating conformally-flat and cylindrical tetrads is given by (details see in Supplement A)

$$\Psi_{cyl} = S \Psi_{cart} , \quad S = \begin{vmatrix} B & 0 \\ 0 & B \end{vmatrix} , \quad (8.3b)$$

$$B = \sigma \begin{vmatrix} A e^{+i\phi/2} & C e^{-i\phi/2} \\ -C e^{+i\phi/2} & A e^{-i\phi/2} \end{vmatrix} , \quad \sigma = \pm 1 ,$$

$$A = \sqrt{\frac{(1 + \cos \rho)}{(1 + \cos \rho \cos z)}} \cos \frac{z}{2} , \quad C = \sqrt{\frac{(1 - \cos \rho)}{(1 + \cos \rho \cos z)}} \sin \frac{z}{2} .$$

One can calculate an explicit form of $\hat{l}_3$ in conformally-flat basis

$$\begin{aligned} \hat{l}_3 = -i\partial_\phi &\implies B^{-1}\hat{l}_3 B = -iB^{-1}\partial_\phi B = \\ &= -i\partial_\phi + \frac{1}{2} \begin{vmatrix} A e^{-i\phi/2} & -C e^{-i\phi/2} \\ C e^{+i\phi/2} & A e^{+i\phi/2} \end{vmatrix} = -i\partial_\phi + \frac{1}{2} \begin{vmatrix} +A e^{+i\phi/2} & -C e^{-i\phi/2} \\ -C e^{+i\phi/2} & -A e^{-i\phi/2} \end{vmatrix} = \\ &= -i\partial_\phi + \frac{1}{2} \begin{vmatrix} A^2 + C^2 & 0 \\ 0 & -A^2 - C^2 \end{vmatrix} = -i\partial_\phi + \frac{1}{2} \begin{vmatrix} 1 & 0 \\ 0 & -1 \end{vmatrix} = -i\partial_\phi + \frac{1}{2} \sigma^3 , \end{aligned}$$

so

$$cyl \implies cart : \quad B^{-1}\hat{l}_3 B = -i\partial_\phi + \frac{1}{2} \sigma^3 = \hat{J}_3 . \quad (8.4)$$

Now let us calculate the form of the wave function in conformally-flat basis:

$$\psi_{cart} = B^{-1}\psi_{cyl} = e^{im\phi} e^{ikz} \begin{vmatrix} A e^{-i\phi/2} f_1(\rho) - C e^{-i\phi/2} f_2(\rho) \\ C e^{+i\phi/2} f_1(\rho) + A e^{+i\phi/2} f_2(\rho) \end{vmatrix} = \begin{vmatrix} \psi_1 \\ \psi_2 \end{vmatrix} ; \quad (8.5a)$$

for ψ_1 and ψ_2 we obtain expressions

$$\begin{aligned} \psi_1 &= e^{i(m-1/2)\phi} \left[f_1(\rho) \sqrt{\frac{1 + \cos \rho}{1 + \cos \rho \cos z}} \cos \frac{z}{2} e^{ikz} - f_2(\rho) \sqrt{\frac{1 - \cos \rho}{1 + \cos \rho \cos z}} \sin \frac{z}{2} e^{ikz} \right] , \\ \psi_2 &= e^{i(m+1/2)\phi} \left[f_1(\rho) \sqrt{\frac{1 - \cos \rho}{1 + \cos \rho \cos z}} \sin \frac{z}{2} e^{ikz} + f_2(\rho) \sqrt{\frac{1 + \cos \rho}{1 + \cos \rho \cos z}} \cos \frac{z}{2} e^{ikz} \right] . \end{aligned} \quad (8.5b)$$

From where, taking into account

$$\cos \frac{z}{2} e^{ikz} = \frac{e^{i(k+1/2)z} + e^{i(k-1/2)z}}{2} , \quad \sin \frac{z}{2} e^{ikz} = \frac{e^{i(k+1/2)z} - e^{i(k-1/2)z}}{2i} ,$$

and expression for f_1, f_2 in terms of G_1, G_2), it follows

$$\psi_1 = \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{-iz}) G_1(\rho) - i \sin \rho e^{-iz} G_2(\rho) \right] ,$$

$$\psi_2 = \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{+iz}) G_2(\rho) - i \sin \rho e^{+iz} G_1(\rho) \right] . \quad (8.5c)$$

From eqs. (8.5c) one can conclude:

only when m and k are half-integer, the constituents ψ_1 and ψ_2 of the fermion wave function in conformally-flat basis may be single-valued and continuous in the spherical space S_3 .

It may be of interest the form of $\hat{N}_3^{cyl} = -i\partial_z$ after transforming to a conformally-flat basis, $\hat{N}_3^{cart} = -iB^{-1}\partial_z B$:

$$\hat{N}_3^{cart} = -i\partial_z - i \begin{vmatrix} 0 & +e^{-i\phi}(A\partial_z C - C\partial_z A) \\ -e^{i\phi}(A\partial_z C - C\partial_z A) & 0 \end{vmatrix} ; \quad (8.6a)$$

from where taking into account

$$A\partial_z C - C\partial_z A = \frac{\sin \rho}{2(1 + \cos \rho \cos z)^2} ,$$

we get

$$\hat{N}_3^{cart} = -i\partial_z + \frac{i \sin \rho}{2(1 + \cos \rho \cos z)^2} \begin{vmatrix} 0 & -e^{-i\phi} \\ +e^{i\phi} & 0 \end{vmatrix} ; \quad (8.6b)$$

With the use of relations

$$x^1 = \frac{\sin \rho \cos \phi}{1 + \cos \rho \cos z} , \quad x^2 = \frac{\sin \rho \sin \phi}{1 + \cos \rho \cos z} , \quad x^3 = \frac{\cos \rho \sin \phi}{1 + \cos \rho \cos z} , \quad f = \frac{1}{1 + \cos \rho \cos z} ,$$

$\hat{N}_3^{cart}$ from (8.6b) will look as

$$\hat{N}_3^{cart} = -i\partial_z + \frac{i}{2} f(x) \begin{vmatrix} 0 & -(x^1 - ix^2) \\ (x^1 + ix^2) & 0 \end{vmatrix} . \quad (8.6c)$$

9. Analysis of the special cases $\lambda = \pm 1/2$ and $\lambda = \pm 3/2$

The above relation between differential equations of first order (7.8) and equations of second order (7.7) has some peculiarities. Indeed, from (7.8) it follows (7.7) only if

$$\lambda + \frac{1}{2} - m + k \neq 0, \quad \lambda + \frac{1}{2} + m - k \neq 0 . \quad (9.1a)$$

In opposite case

$$\lambda + \frac{1}{2} - m + k = 0, \quad \lambda + \frac{1}{2} + m - k = 0 ,$$

which is possible when

$$\lambda = -1/2, \quad m = k , \quad (9.1b)$$

the system (7.8) turns to be a pair of two separated first order equations:

$$\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} - m \frac{\cos \rho}{\sin \rho} - m \frac{\sin \rho}{\cos \rho} \right) G_1 = 0 , \quad (9.2a)$$

$$\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} + m \frac{\cos \rho}{\sin \rho} + m \frac{\sin \rho}{\cos \rho} \right) G_2 = 0 . \quad (9.2b)$$

Eqs. (9.2) may be quite easily solved. Indeed, they can be rewritten as

$$\left(\frac{d}{d\rho} + \frac{\cos 2\rho - 2m}{\sin 2\rho} \right) G_1 = 0 , \quad \left(\frac{d}{d\rho} + \frac{\cos 2\rho + 2m}{\sin 2\rho} \right) G_2 = 0 ,$$

and further

$$\frac{G'_1}{G_1} = + \frac{2m}{\sin 2\rho} - \frac{\cos 2\rho}{\sin 2\rho} , \quad \frac{G'_2}{G_2} = - \frac{2m}{\sin 2\rho} - \frac{\cos 2\rho}{\sin 2\rho} ,$$

these can be readily integrated:

$$\ln G_1 = \ln C_1 + m \ln(\tan \rho) - \frac{1}{2} \ln(\sin 2\rho) ,$$

$$\ln G_2 = \ln C_2 - m \ln(\tan \rho) - \frac{1}{2} \ln(\sin 2\rho) .$$

Thus, the following class of solutions is found:

$$\underline{\lambda = -1/2, \quad k = +m} , \quad G_1(\rho) = C_1 \frac{\tan^m \rho}{\sqrt{\sin 2\rho}} , \quad G_2(\rho) = C_2 \frac{\tan^{-m} \rho}{\sqrt{\sin 2\rho}} . \quad (9.3)$$

Take notice that the constructed wave functions (9.3) are singular in the points $\rho = 0, \pi/2$. This makes necessary some additional analysis because one must exclude all solutions not-finite or discontinuous in the space S_3 .

For instance, let us consider the solution

$$\underline{\lambda = -1/2, \quad k = +1/2, m = +1/2} ,$$

$$G_1(\rho) = \frac{C_1}{\sqrt{2} \cos \rho} , \quad G_2(\rho) = \frac{C_2}{\sqrt{2} \sin \rho} . \quad (9.4a)$$

It may be viewed as physical only if its singularities will be eliminated by means of the gauge transformation involved. In conformally-flat tetrad this solution will read as

$$\psi_1 = \frac{e^{iz}}{\sqrt{2}\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{-iz}) \frac{C_1}{\cos \rho} - i \sin \rho e^{-iz} \frac{C_2}{\sin \rho} \right] ,$$

$$\psi_2 = \frac{e^{i\phi}}{\sqrt{2}\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{+iz}) \frac{C_2}{\sin \rho} - i \sin \rho e^{+iz} \frac{C_1}{\cos \rho} \right] . \quad (9.4b)$$

When $\rho = 0$ we have

$$\psi_1 \rightarrow \frac{e^{iz}}{\sqrt{2}\sqrt{1 + \cos z}} \left[(1 + e^{-iz}) C_1 - i e^{-iz} C_2 \right] ,$$

$$\psi_2 \rightarrow \frac{e^{i\phi}}{\sqrt{2}\sqrt{1+\cos z}} \left[(1+e^{+iz}) \frac{C_2}{0} - 0 i e^{+iz} C_1 \right]. \quad (9.4c)$$

The solution must be considered on the closed curve $n_3 = \sin z$, $n_4 = \cos z$, so to be continuous in на S_3 it may depend only on the z , or it must be a constant. From this it follows $C_2 = 0$. When $\rho = \pi/2$ we have

$$\Psi_1 \rightarrow \frac{e^{iz}}{\sqrt{2}} \left[\frac{C_1}{0} - i e^{-iz} C_2 \right], \quad \Psi_2 \rightarrow \frac{e^{i\phi}}{\sqrt{2}} \left[C_2 - i e^{+iz} \frac{C_1}{0} \right]. \quad (9.4d)$$

Here, the solution must be considered on the closed curve $n_1 = \cos \phi$, $n_2 = \sin \phi$, so to be continuous in на S_3 it may depend only on the ϕ or it must be a constant. From this it follows $C_1 = 0$. Thus, the solution (9.4a)-(9.4b) is a good in the sense of continuity only in trivial case $C_1 = 0, C_2 = 0$. Therefore, this solution must be excluded.

In the same manner one can treat another (symmetrical) case:

$$\underline{\lambda = -1/2}, \quad k = -1/2, m = -1/2, \\ G_1(\rho) = \frac{C_1}{\sqrt{2} \sin \rho}, \quad G_2(\rho) = \frac{C_2}{\sqrt{2} \cos \rho}; \quad (9.5a)$$

in conformally-flat basis it looks as

$$\psi_1 = \frac{e^{-i\phi}}{\sqrt{2}\sqrt{1+\cos \rho \cos z}} \left[(1+\cos \rho e^{-iz}) \frac{C_1}{\sin \rho} - i \sin \rho e^{-iz} \frac{C_2}{\cos \rho} \right], \\ \psi_2 = \frac{e^{-iz}}{\sqrt{2}\sqrt{1+\cos \rho \cos z}} \left[(1+\cos \rho e^{+iz}) \frac{C_2}{\cos \rho} - i \sin \rho e^{+iz} \frac{C_1}{\sin \rho} \right]. \quad (9.5b)$$

When $\rho = 0$

$$\psi_1 \rightarrow \frac{e^{-i\phi}}{\sqrt{2}\sqrt{1+\cos z}} \left[(1+e^{-iz}) \frac{C_1}{0} - i 0 e^{-iz} C_2 \right], \\ \psi_2 \rightarrow \frac{e^{-iz}}{\sqrt{2}\sqrt{1+\cos z}} \left[(1+e^{+iz}) C_2 - i e^{+iz} C_1 \right]. \quad (9.5c)$$

When $\rho = \pi/2$

$$\psi_1 \rightarrow \frac{e^{-i\phi}}{\sqrt{2}} \left[C_1 - i e^{-iz} \frac{C_2}{0} \right], \\ \psi_2 \rightarrow \frac{e^{-iz}}{\sqrt{2}} \left[\frac{C_2}{0} - i e^{+iz} C_1 \right]. \quad (9.5d)$$

Evidently, this solution must be excluded by continuity reason.

All the remaining solutions of the type (9.3)

$$\underline{\lambda = -1/2}, \quad k = N + 1/2, m = N + 1/2, \quad (N = 1, 2, \dots), \\ G_1(\rho) = \frac{\sin^N \rho}{\cos^N \rho} \frac{C_1}{\sqrt{2} \cos \rho}, \quad G_2(\rho) = \frac{\cos^N \rho}{\sin^N \rho} \frac{C_2}{\sqrt{2} \sin \rho};$$

$$\underline{\lambda = -1/2}, \quad k = -N - 1/2, m = -N - 1/2, \quad N = 1, 2, \dots,$$

$$G_1(\rho) = \frac{\cos^N \rho}{\sin^N \rho} \frac{C_1}{\sqrt{2} \sin \rho}, \quad G_2(\rho) = \frac{\sin^N \rho}{\cos^N \rho} \frac{C_2}{\sqrt{2} \cos \rho}.$$

are not finite in all tetrad bases and they must be excluded.

There are two other classes of simplest solutions which should be considered here as well: they. Indeed, let it be

$$G_1(\rho) = C_1 = \text{const}, \quad m = +1/2, \quad k = -1/2. \quad (9.6a)$$

then eq. (7.8) take the form

$$\left(\frac{\cos 2\rho}{\sin 2\rho} - \frac{\cos \rho}{2 \sin \rho} + \frac{\sin \rho}{2 \cos \rho} \right) C_1 = i (\lambda - 1/2) G_2(\rho),$$

$$\left(\frac{d}{d\rho} + \frac{\cos 2\rho}{\sin 2\rho} + \frac{\cos \rho}{2 \sin \rho} - \frac{\sin \rho}{2 \cos \rho} \right) G_2(\rho) = i (\lambda + 3/2) C_1,$$

or

$$\begin{aligned} 0 &= (\lambda - 1/2) G_2(\rho), \\ \left(\frac{d}{d\rho} + 2 \frac{\cos 2\rho}{\sin 2\rho} \right) G_2(\rho) &= i (\lambda + 3/2) C_1, \end{aligned} \quad (9.6b)$$

there can be separated three possibilities:

$$\begin{aligned} (1) \quad \lambda &\neq 1/2, -3/2: & C_1(\rho) &= 0, \quad G_2(\rho) = 0, \\ (2) \quad \lambda &= -3/2: & G_1(\rho) &= C_1, \quad G_2(\rho) = 0, \\ (3) \quad \lambda &= +1/2: & \left(\frac{d}{d\rho} + 2 \frac{\cos 2\rho}{\sin 2\rho} \right) G_2(\rho) &= 2i C_1. \end{aligned} \quad (9.6c)$$

In the last case $\lambda = +1/2$ equations are readily integrated:

$$\begin{aligned} G_2 &= \frac{H(\rho)}{\sin 2\rho}, \quad \left(\frac{d}{d\rho} + 2 \frac{\cos 2\rho}{\sin 2\rho} \right) \frac{H(\rho)}{\sin 2\rho} = \frac{1}{\sin 2\rho} \frac{dH}{d\rho}, \quad \implies \\ \frac{dH}{d\rho} &= 2i C_1 \sin 2\rho, \quad H = C_2 - i C_1 \cos 2\rho, \end{aligned}$$

that is

$$G_2(\rho) = \frac{C_2 - i C_1 \cos 2\rho}{\sin 2\rho}.$$

Thus, two solutiona constaructed:

$$\begin{aligned} (2): \quad \underline{\lambda = -3/2}, \quad m &= +1/2, \quad k = -1/2, \\ G_1(\rho) &= C_1, \quad G_2(\rho) = 0; \\ \Psi_1 &= \frac{1}{\sqrt{1 + \cos \rho \cos z}} (1 + \cos \rho e^{-iz}) C_1, \\ \Psi_2 &= \frac{e^{+i\phi} e^{-iz}}{\sqrt{1 + \cos \rho \cos z}} (-i \sin \rho e^{+iz} C_1); \end{aligned} \quad (9.7)$$

$$\rho = 0 : \quad \Psi_1 = \frac{C_1(1 + e^{-iz})}{\sqrt{1 + \cos z}}, \quad \Psi_2 = 0.$$

$$\rho = \pi/2 : \quad \Psi_1 = C_1, \quad \Psi_2 = -iC_1 e^{+i\phi}.$$

it is finite and continuous in S_3

$$(3) : \quad \underline{\lambda = +1/2}, \quad m = +1/2, \quad k = -1/2,$$

$$G_1(\rho) = C_1, \quad G_2(\rho) = \frac{C_2 - iC_1 \cos 2\rho}{2 \sin \rho \cos \rho}. \quad (9.8a)$$

$$\Psi_1 = \frac{1}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{-iz}) C_1 - i \sin \rho e^{-iz} \frac{C_2 - iC_1 \cos 2\rho}{2 \sin \rho \cos \rho} \right],$$

$$\Psi_2 = \frac{e^{i\phi} e^{-iz}}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{+iz}) \frac{C_2 - iC_1 \cos 2\rho}{2 \sin \rho \cos \rho} - i \sin \rho e^{+iz} C_1 \right]; \quad (9.8b)$$

$$\rho = 0 : \quad \Psi_1 = \frac{1}{\sqrt{1 + \cos z}} \left[(1 + e^{-iz}) C_1 - i e^{-iz} \frac{C_2 - iC_1}{2} \right],$$

$$\rho = 0 : \quad \Psi_2 = \frac{e^{i\phi} e^{-iz}}{\sqrt{1 + \cos z}} (1 + e^{+iz}) \frac{C_2 - iC_1}{0};$$

$$\rho = \pi/2 : \quad \Psi_1 = C_1 - i e^{-iz} \frac{C_2 + iC_1}{0},$$

$$\rho = \pi/2 : \quad \Psi_2 = e^{i\phi} e^{-iz} \left[\frac{C_2 + iC_1}{0} - i e^{+iz} C_1 \right]. \quad (9.8c)$$

it must be rejected. by continuity reason. There exists another symmetrical variant:

$$G_2(\rho) = C_2 = \text{const}, \quad m = -1/2, \quad k = +1/2, \quad (9.9a)$$

then (7.8) takes the form

$$0 = (\lambda - 1/2) G_1(\rho),$$

$$\left(\frac{d}{d\rho} + 2 \frac{\cos 2\rho}{\sin 2\rho} \right) G_1(\rho) = i (\lambda + 3/2) C_2, \quad (9.9b)$$

Depending on three possibilities

$$(1) \quad \lambda \neq +1/2, -3/2 : \quad C_1(\rho) = 0, \quad G_2(\rho) = 0,$$

$$(2) \quad \lambda = -3/2 : \quad G_1(\rho) = 0, \quad G_2(\rho) = C_2, \quad (9.9c)$$

$$(3) \quad \lambda = +1/2 : \quad \left(\frac{d}{d\rho} + 2 \frac{\cos 2\rho}{\sin 2\rho} \right) G_1(\rho) = 2i C_2.$$

we can obtain not trivial solutions. When $\lambda = +1/2$

$$(3) \quad G_1(\rho) = \frac{C_1 - iC_2 \cos 2\rho}{\sin 2\rho}.$$

Thus, two solutions are constructed:

$$(2) \quad \underline{\lambda = -3/2}, \quad m = -1/2, \quad k = +1/2,$$

$$G_1(\rho) = 0, \quad G_2(\rho) = C_2 ; \quad (9.10a)$$

$$(3) \quad \underline{\lambda = +1/2}, \quad m = -1/2, \quad k = +1/2, \\ \lambda = +1/2 : \quad G_2(\rho) = C_2, \quad G_1(\rho) = \frac{C_1 - iC_2 \cos 2\rho}{\sin 2\rho}. \quad (9.10b)$$

Only the solution (2) turns to be finite and continuous in S_3 . Its cartesian form and behavior in special points is given below:

$$(2) \quad \underline{\lambda = -3/2, \quad m = -1/2, \quad k = +1/2}, \\ \psi_1 = \frac{e^{-i\phi} e^{iz}}{\sqrt{1 + \cos \rho \cos z}} (-i \sin \rho e^{-iz} C_2), \\ \psi_2 = \frac{1}{\sqrt{1 + \cos \rho \cos z}} (1 + \cos \rho e^{+iz}) C_2. \quad (9.11) \\ \rho = 0 : \quad \psi_1 = 0, \quad \psi_2 = \frac{C_2(1 + e^{+iz})}{\sqrt{1 + \cos z}}, \\ \rho = \pi/2 : \quad \psi_1 = -iC_2 e^{-i\phi}, \quad \psi_2 = C_2.$$

It is a good function in the sense of continuity.

The general conclusion may be done: no solutions with $\lambda = \pm 1/2$, continuous in S_3 , exist. The Dirac wave functions with $\lambda = \pm 3/2$ are constructed – see (9.7) and (9.11).

10. Quantization of the Dirac energy levels

Let us turn again to the first order equations (7.3)

$$\left(\frac{d}{d\rho} - m \frac{\cos \rho}{\sin \rho} - k \frac{\sin \rho}{\cos \rho} \right) F_1 = i \left(\lambda + \frac{1}{2} - m + k \right) F_2, \quad (10.1a)$$

$$\left(\frac{d}{d\rho} + m \frac{\cos \rho}{\sin \rho} + k \frac{\sin \rho}{\cos \rho} \right) F_2 = i \left(\lambda + \frac{1}{2} + m - k \right) F_1, \quad (10.1b)$$

or equivalent to them second order equations (7.4):

$$\left[\frac{d^2}{d\rho^2} - \frac{m(m-1)}{\sin^2 \rho} - \frac{k(k+1)}{\cos^2 \rho} + \left(\lambda + \frac{1}{2} \right)^2 \right] F_1(\rho) = 0, \quad (10.2a)$$

$$\left[\frac{d^2}{d\rho^2} - \frac{m(m+1)}{\sin^2 \rho} - \frac{k(k-1)}{\cos^2 \rho} + \left(\lambda + \frac{1}{2} \right)^2 \right] F_2(\rho) = 0. \quad (10.2b)$$

The values of energy can be established if all the function involved are polynomials and the quantum number m and n are taken to be half-integer.

The radial functions F_1, F_2 are taken in the form (calculation can be done for one function only)

$$F_1 = \sin^{A_1} \rho \cos^{B_1} \rho R_1(\rho), \quad F_2 = \sin^{A_2} \rho \cos^{B_2} \rho R_2(\rho). \quad (10.3)$$

Substituting (10.3) into (10.2a), one gets

$$\begin{aligned} \frac{d^2}{d\rho^2} R_1 + (2A_1 \frac{\cos \rho}{\sin \rho} - 2B_2 \frac{\sin \rho}{\cos \rho}) \frac{d}{d\rho} R_1 + \left[A_1(A_1 - 1) \frac{\cos^2 \rho}{\sin^2 \rho} - \frac{m(m-1)}{\sin^2 \rho} + \right. \\ \left. + B_1(B_1 - 1) \frac{\sin^2 \rho}{\cos^2 \rho} - \frac{k(k+1)}{\cos^2 \rho} - A_1(B_1 + 1) - B_1(A_1 + 1) + (\lambda + \frac{1}{2})^2 \right] R_1(\rho) = 0. \end{aligned} \quad (10.4)$$

Let A_1, B_1 obey the relations:

$$\begin{aligned} A_1(A_1 - 1) = m(m-1) &\implies A_1 = m, -m+1 = \pm(-m+1/2) + 1/2, \\ B_1(B_1 - 1) = k(k+1) &\implies B_1 = -k, k+1 = \pm(k+1/2) + 1/2. \end{aligned} \quad (10.5)$$

Then eq. (10.4) will be much simpler

$$\frac{d^2}{d\rho^2} R_1 + 2(A_1 \frac{\cos \rho}{\sin \rho} - B_1 \frac{\sin \rho}{\cos \rho}) \frac{d}{d\rho} R_1 + [(\lambda + \frac{1}{2})^2 - (A_1 + B_1)^2] R_1 = 0. \quad (10.6)$$

In the same way, for R_2 we have

$$\begin{aligned} A_2(A_2 - 1) = m(m+1) &\implies A_2 = -m, m+1 = \pm(m+1/2) + 1/2, \\ B_2(B_2 - 1) = k(k-1) &\implies B_2 = k, -k+1 = \pm(-k+1/2) + 1/2. \end{aligned} \quad (10.7)$$

Introducing a new variable $\cos^2 \rho = y$, eq. (10.6) is changed to the form

$$\begin{aligned} y(1-y) \frac{d^2}{dy^2} R_1 + [B_1 + \frac{1}{2} - (A_1 + B_1 + 1)y] \frac{d}{dy} R_1 + \\ + \frac{1}{4} [(\lambda + 1/2)^2 - (A_1 + B_1)^2] R_1 = 0, \end{aligned}$$

that can be compared with the hypergeometric case

$$\begin{aligned} R_1 = C_1 F(\alpha_1, \beta_1, \gamma_1; y), \quad \gamma_1 = B_1 + 1/2 \\ A_1 + B_1 + 1 = \alpha_1 + \beta_1 + 1, \quad \frac{(\lambda + 1/2)^2 - (A_1 + B_1)^2}{4} = -\alpha_1 \beta_1; \end{aligned} \quad (10.8b)$$

with α_1, β_1 taken as

$$\alpha_1 = \frac{A_1 + B_1 + (\lambda + 1/2)}{2}, \quad \beta_1 = \frac{A_1 + B_1 - (\lambda + 1/2)}{2}. \quad (10.8b)$$

Analogously, in the case of R_2 :

$$\begin{aligned} R_2 = C_2 F(\alpha_2, \beta_2, \gamma_2; y), \quad \gamma_2 = B_2 + 1/2, \\ \alpha_2 = \frac{A_2 + B_2 + (\lambda + 1/2)}{2}, \quad \beta_2 = \frac{A_2 + B_2 - (\lambda + 1/2)}{2}. \end{aligned} \quad (10.9)$$

Taking in mind (10.5) and (10.7), instead of A_1, B_1, A_2, B_2 one can introduce

$$A_1 = a_1 + \frac{1}{2}, \quad B_1 = b_1 + \frac{1}{2}, \quad A_2 = a_2 + \frac{1}{2}, \quad B_2 = b_2 + \frac{1}{2}, \quad (10.10)$$

then

$$\begin{aligned}
G_1(\rho) &= C_1 \sin^{a_1} \rho \cos^{b_1} \rho F(\alpha_1, \beta_1, \gamma_1; \cos^2 \rho), \\
G_2(\rho) &= C_2 \sin^{a_2} \rho \cos^{b_2} \rho F(\alpha_2, \beta_2, \gamma_2; \cos^2 \rho), \\
a_1 &= |m - 1/2|, \quad b_1 = |k + 1/2|, \\
a_2 &= |m + 1/2|, \quad b_2 = |k - 1/2|, \\
a_i, b_i &\in \{0, +1, +2, +3, \dots\}.
\end{aligned} \tag{10.11a}$$

Parameters in hypergeometric functions (10.11a) are defined by

$$\begin{aligned}
\lambda &= \pm \sqrt{\epsilon^2 - M^2}, \\
\alpha_1 &= \frac{a_1 + b_1 + 1 + (\lambda + 1/2)}{2}, \quad \beta_1 = \frac{a_1 + b_1 + 1 - (\lambda + 1/2)}{2}, \quad \gamma_1 = b_1 + 1, \\
\alpha_2 &= \frac{a_2 + b_2 + 1 + (\lambda + 1/2)}{2}, \quad \beta_2 = \frac{a_2 + b_2 + 1 - (\lambda + 1/2)}{2}, \quad \gamma_2 = b_2 + 1.
\end{aligned} \tag{10.11b}$$

Depending on the sign of λ the hypergeometric series are the polynomials according to:

$$\begin{aligned}
\lambda > 0, \quad \beta_1 &= \frac{a_1 + b_1 + 1 - (\lambda + 1/2)}{2} = -n_1, \quad \beta_2 = \frac{a_2 + b_2 + 1 - (\lambda + 1/2)}{2} = -n_2, \\
\lambda < 0, \quad \alpha_1 &= \frac{a_1 + b_1 + 1 + (\lambda + 1/2)}{2} = -n_1, \quad \alpha_2 = \frac{a_2 + b_2 + 1 + (\lambda + 1/2)}{2} = -n_2.
\end{aligned} \tag{10.12}$$

Evidently, in view of existence of first order differential relation between F_1 and F_2 one can examine only one second order equation, wether for F_1 or for F_2 .

From the second order equation for $G_1(\rho)$ it follows the quantization rule:

$$\begin{aligned}
\lambda &= +3/2, +5/2, +7/2, \dots, \\
\beta_1 &= \frac{a_1 + b_1 + 1 - (\lambda + 1/2)}{2} = -n_1,
\end{aligned} \tag{10.13a}$$

$$\begin{aligned}
\lambda &= +(a_1 + b_1 + 1/2 + 2n_1), \quad \epsilon^2 = M^2 + \lambda^2, \\
G_1(\rho) &= C_1 \sin^{a_1} \rho \cos^{b_1} \rho F(a_1 + b_1 + 1 + n_1, -n_1, b_1 + 1; \cos^2 \rho) \\
\lambda &= -3/2, -5/2, -7/2, \dots,
\end{aligned}$$

$$\alpha_1 = \frac{a_1 + b_1 + 1 + (\lambda + 1/2)}{2} = -n_1, \tag{10.13b}$$

$$\begin{aligned}
\lambda &= -(a_1 + b_1 + 3/2 + 2n_1), \quad \epsilon^2 = M^2 + \lambda^2, \\
G_1(\rho) &= C_1 \sin^{a_1} \rho \cos^{b_1} \rho F(-n_1, a_1 + b_1 + 1 + n_1, b_1 + 1; \cos^2 \rho)
\end{aligned}$$

All the energy levels are degenerated in λ, m, k :

$$\Psi_\epsilon = \Psi_{\epsilon, \lambda, m, k}, \quad \epsilon = \sqrt{M^2 + \lambda^2},$$

$$\lambda = \pm 3/2, \pm 5/2, \pm 7/2, \dots \quad (10.14)$$

Quantization of λ is given by

$$\lambda = a_1 + b_1 + 1/2 + 2n_1 \quad \text{or} \quad -\lambda = a_1 + b_1 + 3/2 + 2n_1 . \quad (10.15)$$

Some first energy levels may be readily detailed:

$$\begin{aligned}
\underline{\lambda = +3/2} : \quad 1 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (1, 0), (0, 1) . \\
\underline{\lambda = -3/2} : \quad 0 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (0, 0) . \\
\underline{\lambda = +5/2} : \quad 2 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (0, 2), (1, 1), (2, 0) , \\
&n_1 = 1 , \quad (a_1, b_1) = (0, 0) . \\
\underline{\lambda = -5/2} : \quad 1 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (1, 0), (0, 1) . \\
\underline{\lambda = +7/2} : \quad 3 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (0, 3), (1, 2), (2, 1), (3, 0) , \\
&n_1 = 1 , \quad (a_1, b_1) = (0, 1), (1, 0) . \\
\underline{\lambda = -7/2} : \quad 2 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (0, 2), (1, 1), (2, 0) , \\
&n_1 = 1 , \quad (a_1, b_1) = (0, 0) . \\
\underline{\lambda = +9/2} : \quad 4 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (0, 4), (1, 3), (3, 1), (4, 0) , \\
&n_1 = 1 , \quad (a_1, b_1) = (0, 2), (1, 1), (2, 0) , \\
&n_1 = 2 , \quad (a_1, b_1) = (0, 0) . \\
\underline{\lambda = -9/2} : \quad 3 &= a_1 + b_1 + 2n_1 \quad \implies \\
&n_1 = 0 , \quad (a_1, b_1) = (0, 3), (1, 2), (2, 1), (3, 0) , \\
&n_2 = 1, (a_1, b_1) = (0, 1), (1, 0) .
\end{aligned}$$

and so on.

11. The wave functions constructed and continuity testing

All physical Dirac wave solutions must be single-valued and continuous in S_3 . Solutions with minimal values of energy

$$\epsilon = \sqrt{M^2 + (3/2)^2}, \quad \lambda = \pm 3/2 \quad (11.1a)$$

are just those (see §9). It remains to consider solutions with

$$\epsilon = \sqrt{M^2 + \lambda^2}, \quad |\lambda| = 5/2, 7/2, 9/2, \dots \quad (11.1b)$$

Half-integer values for m, k will guarantee continuity of the solutions when $0 < \rho < \pi/2$. Special cases are two closed curves:

$$\rho = 0 : \quad n_3 = \sin z, \quad n_4 = \cos z, \quad (11.2a)$$

$$\rho = \pi/2 : \quad n_1 = \cos \phi, \quad n_2 = \sin \phi. \quad (11.2b)$$

In conformally flat tetrad the wave functions must behave themselves in accordance with the rules

$$\rho = 0 : \quad \Psi_{cart} = \{\psi(z), \text{ or } \text{const}\}, \quad (11.2a)$$

$$\rho = \pi/2 : \quad \Psi_{cart} = \{\psi(\phi), \text{ or } \text{const}\}. \quad (11.2b)$$

Two first components of the wave functions (see (8.5c))

$$\begin{aligned} \psi_1 &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{-iz}) G_1(\rho) - i \sin \rho e^{-iz} G_2(\rho) \right], \\ \psi_2 &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{+iz}) G_2(\rho) - i \sin \rho e^{+iz} G_1(\rho) \right] \end{aligned}$$

will take the form

$$\begin{aligned} \rho = 0 : \quad \psi_1 &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos z}} (1 + e^{-iz}) G_1(0), \\ \rho = 0 : \quad \psi_2 &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos z}} (1 + e^{+iz}) G_2(0), \\ \rho = \pi/2 : \quad \psi_1 &= e^{i(m-1/2)\phi} e^{i(k+1/2)z} \left[G_1(\pi/2) - i e^{-iz} G_2(\pi/2) \right], \\ \rho = \pi/2 : \quad \psi_2 &= e^{i(m+1/2)\phi} e^{i(k-1/2)z} \left[G_2(\pi/2) - i e^{+iz} G_1(\pi/2) \right] \end{aligned} \quad (11.3)$$

where

$$\begin{aligned} G_1(\rho) &= C_1 \sin^{a_1} \rho \cos^{b_1} \rho F(\alpha_1, \beta_1, \gamma_1; \cos^2 \rho), \\ G_2(\rho) &= C_2 \sin^{a_2} \rho \cos^{b_2} \rho F(\alpha_2, \beta_2, \gamma_2; \cos^2 \rho), \\ a_1 &= |m - 1/2|, \quad b_1 = |k + 1/2|, \quad a_2 = |m + 1/2|, \quad b_2 = |k - 1/2|. \end{aligned} \quad (11.4)$$

While all four parameters $a_1, b_1, a_2, b_2 \neq 0$, because of the presence of the terms $\sin^{a_1} \rho \cos^{b_1}$ and $\sin^{a_2} \rho \cos^{b_2} \rho$ the functions $G_1(\rho), G_2(\rho)$ will be equal to zero when $\rho = 0, \pi/2$. Correspondingly, eqs. (11.3) read

$$\begin{aligned} \rho = 0 : \quad \psi_1 &= 0, \quad \psi_2 = 0, \\ \rho = \pi/2 : \quad \psi_1 &= 0, \quad \psi_2 = 0. \end{aligned}$$

All the remaining cases

$$\begin{aligned} m = +1/2, k = +1/2, & \quad m = -1/2, k = -1/2, \\ m = +1/2, k = -1/2, & \quad m = -1/2, k = +1/2, \end{aligned} \quad (11.5a)$$

$$\begin{aligned} m = +1/2, |k| > 1/2, & \quad m = -1/2, |k| > 1/2, \\ |m| > 1/2, k = -1/2, & \quad |m| > 1/2, k = +1/2, \end{aligned} \quad (11.5b)$$

must be considered separately. First, let us turn to the four cases in (11.5a):

$m = +1/2, k = +1/2$:

$$\begin{aligned} a_1 &= 0, b_1 = 1, a_2 = 1, b_2 = 0, \\ G_1(0) &= \text{const}, \quad G_2(0) = 0, \\ G_1(\pi/2) &= 0, \quad G_2(\pi/2) = \text{const} \\ \rho = 0 : \quad \psi_1 &= \frac{e^{iz}(1 + e^{-iz})}{\sqrt{1 + \cos z}} \text{const} = \psi(z), \\ \rho = 0 : \quad \psi_2 &= \frac{e^{i\phi}(1 + e^{+iz})}{\sqrt{1 + \cos z}} 0 = 0. \\ \rho = \pi/2 : \quad \psi_1 &= e^{iz} (0 - i e^{-iz} \text{const}) = -i \text{const}, \\ \rho = \pi/2 : \quad \psi_2 &= e^{i\phi} (\text{const} - i e^{+iz} 0) = \text{const} e^{i\phi}. \end{aligned} \quad (11.6)$$

$m = -1/2, k = -1/2$:

$$\begin{aligned} a_1 &= 1, b_1 = 0, a_2 = 0, b_2 = 1, \\ G_1(0) &= 0, \quad G_2(0) = \text{const}, \\ G_1(\pi/2) &= \text{const}, \quad G_2(\pi/2) = 0 \\ \rho = 0 : \quad \psi_1 &= \frac{e^{-i\phi}}{\sqrt{1 + \cos z}} (1 + e^{-iz}) 0 = 0, \\ \rho = 0 : \quad \psi_2 &= \frac{e^{-iz}(1 + e^{+iz})}{\sqrt{1 + \cos z}} \text{const}. \\ \rho = \pi/2 : \quad \psi_1 &= e^{-i\phi} (\text{const} - i e^{-iz} 0) = \text{const} e^{-i\phi}, \\ \rho = \pi/2 : \quad \psi_2 &= e^{-iz} (0 - i e^{+iz} \text{const}) = -i \text{const}. \end{aligned} \quad (11.7)$$

$m = +1/2, k = -1/2$:

$$\begin{aligned} a_1 &= 0, b_1 = 0, a_2 = 1, b_2 = 1, \\ G_1(0) &= \text{const}, \quad G_2(0) = 0, \\ G_1(\pi/2) &= \text{const}, \quad G_2(\pi/2) = 0 \\ \rho = 0 : \quad \psi_1 &= \frac{(1 + e^{-iz})}{\sqrt{1 + \cos z}} \text{const}, \end{aligned}$$

$$\rho = 0 : \quad \psi_2 = \frac{e^{i\phi} e^{-iz} (1 + e^{+iz})}{\sqrt{1 + \cos z}} 0 = 0 . \quad (11.8)$$

$$\rho = \pi/2 : \quad \psi_1 = (\text{const} - i e^{-iz} 0) = \text{const} ,$$

$$\rho = \pi/2 : \quad \psi_2 = e^{i\phi} e^{-iz} (0 - i e^{+iz} \text{const}) = -i e^{i\phi} .$$

$m = -1/2, k = +1/2$:

$$a_1 = 1 , \quad b_1 = 1 , \quad a_2 = 0 , \quad b_2 = 0 ,$$

$$G_1(0) = 0 , \quad G_2(0) = \text{const} ,$$

$$G_1(\pi/2) = 0 , \quad G_2(\pi/2) = \text{const} ,$$

$$\rho = 0 : \quad \psi_1 = \frac{e^{-i\phi} e^{iz} (1 + e^{-iz})}{\sqrt{1 + \cos z}} 0 = 0 ,$$

$$\rho = 0 : \quad \psi_2 = \frac{(1 + e^{+iz})}{\sqrt{1 + \cos z}} \text{const} . \quad (11.9)$$

$$\rho = \pi/2 : \quad \psi_1 = e^{-i\phi} e^{iz} (0 - i e^{-iz} \text{const}) = -i \text{const} e^{-i\phi} ,$$

$$\rho = \pi/2 : \quad \psi_2 = \text{const} - i e^{+iz} 0 = \text{const} .$$

Now let us consider the cases in (11.5b).

$m = +1/2 , |k| > +1/2$:

$$a_1 = 0 , \quad b_1 \neq 0 , \quad a_2 = 1 , \quad b_2 \neq 0 ,$$

$$G_1(0) = \text{const} , \quad G_2(0) = 0 ,$$

$$G_1(\pi/2) = 0 , \quad G_2(\pi/2) = 0 ,$$

$$\rho = 0 : \quad \psi_1 = \frac{e^{i(k+1/2)z}}{\sqrt{1 + \cos z}} (1 + e^{-iz}) \text{const} = \psi(z) ,$$

$$\rho = 0 : \quad \psi_2 = \frac{e^{i\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos z}} (1 + e^{+iz}) 0 ,$$

$$\rho = \pi/2 : \quad \psi_1 = e^{i(k+1/2)z} [0 - i e^{-iz} 0] = 0 ,$$

$$\rho = \pi/2 : \quad \psi_2 = e^{i\phi} e^{i(k-1/2)z} [0 - i e^{+iz} 0] = 0 . \quad (11.10)$$

$m = -1/2 , |k| > +1/2$:

$$a_1 = 1 , \quad b_1 \neq 0 , \quad a_2 = 0 , \quad b_2 \neq 0 ,$$

$$G_1(0) = 0 , \quad G_2(0) = \text{const} ,$$

$$G_1(\pi/2) = 0 , \quad G_2(\pi/2) = 0 ,$$

$$\rho = 0 : \quad \psi_1 = \frac{e^{-i\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos z}} (1 + e^{-iz}) 0 = 0 ,$$

$$\rho = 0 : \quad \psi_2 = \frac{e^{i(k-1/2)z}}{\sqrt{1 + \cos z}} (1 + e^{+iz}) \text{const} = \psi_2(z) ,$$

$$\rho = \pi/2 : \quad \psi_1 = e^{i(k+1/2)z} [0 - i e^{-iz} 0] = 0 ,$$

$$\rho = \pi/2 : \quad \psi_2 = e^{i\phi} e^{i(k-1/2)z} \begin{bmatrix} 0 - i e^{+iz} & 0 \end{bmatrix} = 0 . \quad (11.11)$$

$|m| > 1/2, k = +1/2:$

$$a_1 \neq 0, b_1 = 1, a_2 \neq 0, b_2 = 0,$$

$$G_1(0) = 0, G_2(0) = 0,$$

$$G_1(\pi/2) = 0, G_2(\pi/2) = \text{const},$$

$$\rho = 0 : \quad \psi_1 = \frac{e^{i(m-1/2)\phi} e^{iz}}{\sqrt{1 + \cos z}} (1 + e^{-iz}) = 0,$$

$$\rho = 0 : \quad \psi_2 = \frac{e^{i(m+1/2)\phi}}{\sqrt{1 + \cos z}} (1 + e^{+iz}) = 0,$$

$$\rho = \pi/2 : \quad \psi_1 = e^{i(m-1/2)\phi} e^{iz} \begin{bmatrix} 0 - i e^{-iz} \text{const} \end{bmatrix} = \psi_1(\phi),$$

$$\rho = \pi/2 : \quad \psi_2 = e^{i(m+1/2)\phi} \begin{bmatrix} \text{const} - i e^{+iz} & 0 \end{bmatrix} = \psi_2(\phi). \quad (11.12)$$

$|m| > 1/2, k = -1/2:$

$$a_1 \neq 0, b_1 = 0, a_2 \neq 0, b_2 = 1,$$

$$G_1(0) = 0, G_2(0) = 0,$$

$$G_1(\pi/2) = \text{const}, G_2(\pi/2) = 0,$$

$$\rho = 0 : \quad \psi_1 = \frac{e^{i(m-1/2)\phi}}{\sqrt{1 + \cos z}} (1 + e^{-iz}) = 0,$$

$$\rho = 0 : \quad \psi_2 = \frac{e^{i(m+1/2)\phi} e^{-iz}}{\sqrt{1 + \cos z}} (1 + e^{+iz}) = 0,$$

$$\rho = \pi/2 : \quad \psi_1 = e^{i(m-1/2)\phi} \begin{bmatrix} \text{const} - i e^{-iz} & 0 \end{bmatrix} = \psi_1(\phi),$$

$$\rho = \pi/2 : \quad \psi_2 = e^{i(m+1/2)\phi} e^{-iz} \begin{bmatrix} 0 - i e^{+iz} \text{const} \end{bmatrix} = \psi_2(\phi). \quad (11.13)$$

Thus, the general conclusion can be done: all the constructed Dirac wave functions in cylindrical coordinates $\Psi_{\epsilon, \lambda, m, k}(t, \rho, \phi, z)$ provide us with single-valued and continuous functions on the sphere S_3 . For proving these properties an explicit form of the wave function in conformally flat tetrad basis has been used.

PART III. THE DIRAC EQUATION IN THE ELLIPTICAL SPACE $\tilde{S}_3$

12. Cylindrical coordinates and tetrad in elliptical space, spinor gauge transformations

Now we are to examine the properties of a fermion particle in elliptical space model $\tilde{S}_3$. As a first step, some analogue of the cartesian tetrad basis in $\tilde{S}_3$ should be defined and the explicit form of Dirac solutions in that basis must be calculated. After that one can investigate such wave solution from the point of view of their continuity in the elliptical manifold $\tilde{S}_3$.

Like in §3 we will employ realizations of spherical and elliptical models as group manifolds $SU(2)$ and $SO(3.R)$ respectively:

$$B = n_0 - i n_j \sigma_j , \quad n_0^2 + n_1^2 + n_2^2 + n_3^2 = +1 , \quad (12.1)$$

$$O(\vec{c}) = I + 2 \frac{\vec{c}^\times + (\vec{c}^\times)^2}{1 + \vec{c}^2} , c^i = \frac{+n_i}{+n_4} = \frac{-n_i}{-n_4} . \quad (12.2)$$

To pair of vectors $\vec{c}^\pm = + \pm \infty \vec{c}_0$, $\vec{c}_0^2 = 1$ corresponds one the same matrix, point in $\tilde{S}_3$: $O(\vec{c}^\pm) = I + 2(\vec{c}_0^\times)^2$. From formal point of view, elliptical model can be obtained by identification of opposite point on 4-sphere. However, such a simple view may be insufficient in applications. Below, a more detailed treatment of this relating procedure will be done. In the following, the coordinates (c^1, c^2, c^3) will be considered as Cartesian ones in the model $\tilde{S}_3$. Let us start with the known formulas for the spherical model:

$$\begin{aligned} x^1 &= \frac{n_1}{1 + n_0} , \quad x^2 = \frac{n_2}{1 + n_0} , \quad x^3 = \frac{n_3}{1 + n_0} , \\ n_i &= \frac{2x^i}{1 + x^2} , \quad n_0 = \frac{1 - x^2}{1 + x^2} , \\ dS^2 &= dt^2 - dl^2(x), \quad dl^2(x) = \frac{1}{f^2} [(dx^1)^2 + (dx^2)^2 + (dx^3)^2] , \\ f &= \frac{1 + x^2}{2} , \quad e_{(a)\alpha}(x) = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & f & 0 & 0 \\ 0 & 0 & f & 0 \\ 0 & 0 & 0 & f \end{vmatrix} . \end{aligned} \quad (12.3a)$$

Let $n_0 = (-1 + \delta)$, where δ stands for a infinitely small quantity , then

$$n_a = (-1 + \delta, 0, 0, 0) \implies \lim_{\delta \rightarrow 0} x^i = \frac{1}{\delta} n_i = \infty n_i . \quad (12.3b)$$

This means that all the set of finite coordinates $\{ x^i = \infty n_i ; \vec{n}^2 = 1 \}$ represents one the same point $n_a = (-1; 0, 0, 0)$ on the sphere. Cartesian c^i -coordinate in $\tilde{S}_3$ can be connected with x^i -coordinates in the S_3 as follows (it is a $2 \implies 1$ mapping):

$$\frac{2x^i}{1 - x^2} = c^i ; \quad (12.4a)$$

to the sets x^i and $x'^i = -x^i/x^2$ corresponds one the same vector $c^i = c'^i$. The inverse map is 2-valued:

$$\delta = \pm 1, \quad x_{(\delta)}^i = \frac{c^i}{(1 + \delta \sqrt{1 + c^2})} ; \quad (12.4b)$$

one should note the identities

$$\begin{aligned} \frac{1 - x_{(\delta)}^2}{2} &= \frac{1}{1 + \delta \sqrt{1 + c^2}} , \quad \frac{1 + x_{(\delta)}^2}{2} = \frac{\delta \sqrt{1 + c^2}}{1 + \delta \sqrt{1 + c^2}} , \\ x_{(+)}^2 &< 1 , \quad x_{(-)}^2 > 1 , \quad (c^2 \rightarrow \infty, x_{(\pm)}^2 \rightarrow 1 \mp 0) , \\ \frac{1}{\sqrt{1 + c^2}} &= \delta \frac{1 - x_{(\delta)}^2}{1 + x_{(\delta)}^2} . \end{aligned} \quad (12.4c)$$

Thus, the whole sphere, composed of two parts, can be covered by two maps:

$$n_{0(\delta)} = \frac{\delta}{\sqrt{1+c^2}} : \quad \implies \quad \begin{aligned} n_{0(+1)} &\in [0, +1] , \\ n_{0(-1)} &\in [-1, 0] , \end{aligned}$$

$$\vec{n}_{(\delta)} = \frac{2\vec{x}_{(\delta)}}{1+x_{(\delta)}^2} = \delta \frac{\vec{c}}{\sqrt{1+c^2}} ; \quad (12.4d)$$

it is convenient to have their form in two limits:

$$\begin{aligned} \vec{c} \rightarrow 0 \vec{c}_0 : \quad & n_{0(+1)} \rightarrow +1 , \quad & \vec{n}_{(+1)} \rightarrow 0 \text{ (+1) } \vec{c}_0 = 0 , \\ \vec{c} \rightarrow \infty \vec{c}_0 : \quad & n_{0(+1)} \rightarrow +1/\infty = 0 , \quad & \vec{n}_{(+1)} \rightarrow + \vec{c}_0 , \\ \vec{c} \rightarrow 0 \vec{c}_0 : \quad & n_{0(-1)} \rightarrow -1 , \quad & \vec{n}_{(-1)} \rightarrow 0 \text{ (-1) } \vec{c}_0 = 0 , \\ \vec{c} \rightarrow \infty \vec{c}_0 : \quad & n_{0(-1)} \rightarrow -1/\infty = 0 , \quad & \vec{n}_{(-1)} \rightarrow - \vec{c}_0 . \end{aligned} \quad (12.4e)$$

So, the mapping with $\delta = +1$ to the zero vector $\vec{c} = (0,0,0)$ refers the point $n_a = (+1, 0, 0, 0)$; whereas the mapping with $\delta = -1$ to the zero vector corresponds the point $n_a = (-1, 0, 0, 0)$.

Now with the use of tensor transformation law let us calculate the metric tensor c_i -coordinates of elliptical space Here we face some peculiarities steaming from $2 \implies 1$ character of the mapping from S_3 to $\tilde{S}_3$. Therefore, we must handle every half-space separately:

$$\begin{aligned} dc^i &= \frac{\partial c^i}{\partial x^{j(\delta)}} dx^{j(\delta)} = \\ &= \left[\frac{2\delta_{ij}}{1-x_{(\delta)}^2} + c^i c^j \right] dx^{j(\delta)} = \left[\delta_{ij} (1 + \delta\sqrt{1+c^2}) + c^i c^j \right] dx^{j(\delta)} , \quad (12.5a) \\ dc^i dc^i &= \left[\frac{2\delta_{ij}}{1-x_{(\delta)}^2} + c^i c^j \right] dx^{j(\delta)} \left[\frac{2\delta_{ik}}{1-x_{(\delta)}^2} + c^i c^k \right] dx^{k(\delta)} = \\ &= \left[\frac{4\delta_{jk}}{(1-x_{(\delta)}^2)^2} + \frac{4c^j c^k}{1-x_{(\delta)}^2} + c^2 c^j c^k \right] dx^{j(\delta)} dx^{k(\delta)} . \\ c^2 &= \frac{4x_{(\delta)}^2}{(1-x_{(\delta)}^2)^2} , \quad dc^i dc^i = \frac{4}{(1-x_{(\delta)}^2)^2} [\delta_{jk} + c^j c^k] dx^{j(\delta)} dx^{k(\delta)} . \end{aligned}$$

So one gets

$$\frac{dc^i dc^i}{1+c^2} = \frac{4}{(1+x_{(\delta)}^2)^2} [\delta_{jk} + c^j c^k] dx^{j(\delta)} dx^{k(\delta)} . \quad (12.5b)$$

In the same way, one obtains

$$\frac{c^j c^k dc^j dc^k}{(1+c^2)^2} = \frac{4}{(1+x_{(\delta)}^2)^2} c^j c^k dx^{j(\delta)} dx^{k(\delta)} . \quad (12.5c)$$

Combining two last relations, we get to

$$dl^2(x_{(\delta)}) = \frac{4 dx^{j(\delta)} dx^{k(\delta)}}{[1+x_{(\delta)}^2]^2} = \left[\frac{\delta_{jk}}{1+c^2} - \frac{c^j c^k}{(1+c^2)^2} \right] dc^j dc^k = dl^2(c) . \quad (12.5d)$$

In the following we will need the contra-variant metrical tensor:

$$g^{kj}(c) = (1 + c^2) (\delta^{kj} + c^k c^j) . \quad (12.5e)$$

When the metric $g_{ik}(c)$ is employed one need to fix a certain tetrad – by definition it obeys

$$e_{(i)j}(c) e_{(i)k}(c) = g_{jk}(c) , \quad g_{jk}(c) = \frac{\delta_{jk}}{1 + c^2} - \frac{c^j c^k}{(1 + c^2)^2} \quad (12.6a)$$

to be satisfied. There exists one method to fix a tetrad, preserving some consistency with the situation in spherical model. It consists in coordinate transforming the conformally flat tetrad $e_{(i)k}(x_{(\delta)})$ (the domains $x_{(+1)}^i$ and $x_{(-1)}^i$ must be treated separately) to c^i -coordinates of the elliptical model:

$$e_{(i)k}(x_{(\delta)}) = \frac{1}{f(x_{(\delta)})} \delta_{ik} , \quad f = \frac{1 + x_{(\delta)}^2}{2} ,$$

$$e_{(i)j}^{(\delta)}(c) = \frac{\partial x_{(\delta)}^k}{\partial c^j} e_{(i)k}(x) = \frac{1}{f(x_{(\delta)})} \frac{\partial x_{(\delta)}^i}{\partial c^j} = \frac{1 + \delta \sqrt{1 + c^2}}{\delta \sqrt{1 + c^2}} \frac{\partial x_{(\delta)}^i}{\partial c^j} . \quad (12.6b)$$

In this way we are able to construct two tetrads satisfying eq. (12.6). The problem is reduced to rather simple calculating:

$$\frac{\partial x_{(\delta)}^i}{\partial c^j} = \frac{\partial}{\partial c^j} \frac{c^i}{1 + \delta \sqrt{1 + c^2}} = \frac{\delta_{ij}}{1 + \delta \sqrt{1 + c^2}} - \frac{\delta c^i c^j}{\sqrt{1 + c^2} (1 + \delta \sqrt{1 + c^2})^2} , \quad (12.6c)$$

and further

$$e_{(i)j}^{(\delta)}(c) = \frac{\delta_{ij}}{\delta \sqrt{1 + c^2}} - \frac{c^i c^j}{(1 + c^2) (1 + \delta \sqrt{1 + c^2})} . \quad (12.6d)$$

These two tetrads can be written down as follows

$$e_{(i)j}^{(\delta)}(c) = \begin{vmatrix} a + b c^1 c^1 & b c^1 c^2 & b c^1 c^3 \\ b c^1 c^2 & a + b c^2 c^2 & b c^2 c^3 \\ b c^1 c^3 & b c^2 c^3 & a + b c^3 c^3 \end{vmatrix} ,$$

$$a = \frac{\delta}{\sqrt{1 + c^2}} , \quad b = - \frac{1}{1 + c^2} \frac{1}{1 + \delta \sqrt{1 + c^2}} . \quad (12.7a)$$

One may perform independent calculation which will lead us to the same result. Indeed, let us look for a solution to tetrad equation in the form

$$e_{i(j)}(c) e_{(j)k}(c) = g_{ik}(c) , \quad e_{i(j)}(c) = A \delta_{ij} + B c^i c^j ,$$

from where it follow two equations

$$A^2 = \frac{1}{1 + c^2} , \quad 2AB + B^2 c^2 = - \frac{1}{(1 + c^2)^2} .$$

that is

$$A = - \frac{\mu}{\sqrt{1 + c^2}} , \quad \mu = \mp 1 , \quad B = \frac{1}{c^2(1 + c^2)} (\mu \sqrt{1 + c^2} + \nu) .$$

With the use of identity

$$\frac{(\mu\sqrt{1+c^2} + \nu)}{c^2} = \frac{1}{(\mu\sqrt{1+c^2} - \nu)} ,$$

B can be taken to the form

$$B = \frac{1}{1+c^2} \frac{1}{(\mu\sqrt{1+c^2} - \nu)} = -\nu \frac{1}{1+c^2} \frac{1}{(1 - \nu\mu\sqrt{1+c^2})} .$$

Thus, four different variants are possible:

$$\begin{aligned} (1) \quad (\mu, \nu) = (+1, +1) : \quad A &= -\frac{1}{\sqrt{1+c^2}} , \quad B = -\frac{1}{1+c^2} \frac{1}{(1-\sqrt{1+c^2})} , \\ (2) \quad (\mu, \nu) = (-1, +1) : \quad A &= +\frac{1}{\sqrt{1+c^2}} , \quad B = -\frac{1}{1+c^2} \frac{1}{(1+\sqrt{1+c^2})} , \\ (3) \quad (\mu, \nu) = (-1, -1) : \quad A &= +\frac{1}{\sqrt{1+c^2}} , \quad B = \frac{1}{1+c^2} \frac{1}{(1-\sqrt{1+c^2})} , \\ (4) \quad (\mu, \nu) = (+1, -1) : \quad A &= -\frac{1}{\sqrt{1+c^2}} , \quad B = \frac{1}{1+c^2} \frac{1}{(1+\sqrt{1+c^2})} . \end{aligned} \quad (12.7b)$$

The variants (1) and (2) coincide with listed above (12.7a). variants (3) and (4) correspond to tetrads, P -reflected to (1) and (2). These additional possibilities can be omitted.

These tetrads are substantially different which become evident on looking in their form in limiting points:

$$\vec{c} \rightarrow \epsilon(\vec{c}^0) , \quad \vec{c} \rightarrow \pm \epsilon^{-1}(\vec{c}^0) , \quad \epsilon \rightarrow 0 .$$

Indeed, the metric tensor being $g_{ik}(0) = \delta_{ik}$, $g_{ik}(\infty) = \epsilon^2 (\delta_{ij} - c_i^0 c_j^0) \rightarrow 0$ whereas tetrads behave in accordance with

$$\begin{aligned} e_{(i)j}^{(+1)}(0) &= \delta_{ij} , \quad e_{(i)j}^{(+1)}(\infty) = \begin{vmatrix} 1 & 0 \\ 0 & \pm\epsilon(\delta_{ij} - c_i^0 c_j^0) \end{vmatrix} , \\ e_{(i)j}^{(-1)}(0) &= \begin{vmatrix} 1 & 0 \\ 0 & -\delta_{ij} + 2c_i^0 c_j^0 \end{vmatrix} , \quad e_{\alpha}^{(-1)(a)}(\infty) = \begin{vmatrix} 1 & 0 \\ 0 & \mp\epsilon(\delta_{ij} - c_i^0 c_j^0) \end{vmatrix} . \end{aligned}$$

On general considerations, two tetrads (12.7a) must be connected by a local 3-rotation

$$e_{(i)k}^{(-1)}(c) = O_{ij} e_{(j)k}^{(+1)}(c) , \quad (12.7)$$

which can be calculated explicitly (the notation $\sqrt{1+c^2} = s$ is used):

$$\begin{aligned} O_k^l(c) &= O_k^l(c) = e_{(k)i}^{(-1)}(c) g^{ij}(c) e_{j(l)}^{(+1)}(c) = \\ &= \frac{1}{s^2} \left(-s \delta_{ki} - \frac{c^k c^i}{1-s} \right) (\delta_{ij} + c^i c^j) \left(s \delta_{jl} - \frac{c^j c^l}{1+s} \right) = \\ &= -\delta_{kl} + 2 \frac{c^k c^l}{c^2} = -\delta_{kl} + 2c_0^k c_0^l , \quad \vec{c}_0^2 = 1 . \end{aligned} \quad (12.8a)$$

It is a local gauge rotation of the above special type (12.2b):

$$O(\pm\infty \vec{c}_0) = I + 2(\vec{c}_0^\times)^2 = \quad (12.8b)$$

$$= \begin{vmatrix} 1 - 2(c_0^2 c_0^2 + c_0^3 c_0^3) & 2c_0^1 c_0^2 & 2c_0^1 c_0^3 \\ 2c_0^1 c_0^2 & 1 - 2(c_0^1 c_0^1 + c_0^3 c_0^3) & 2c_0^2 c_0^3 \\ 2c_0^3 c_0^1 & 2c_0^3 c_0^2 & 1 - 2(c_0^1 c_0^1 + c_0^2 c_0^2) \end{vmatrix} = -\delta_{kl} + 2c_0^k c_0^l .$$

Now in the elliptical space let us introduce cylindrical coordinates and tetrad. Two half-space of spherical space are given by


$$\begin{aligned} n_0 &= \cos \rho \cos z , \quad n_3 = \cos \rho \sin z , \quad n_1 = \sin \rho \cos \phi , \quad n_2 = \sin \rho \sin \phi , \\ \underline{\delta = +1}, \quad n_0 &= \cos \rho \cos z \in [0, +1] , \quad z \in [-\frac{\pi}{2}, \frac{\pi}{2}] ; \\ \underline{\delta = -1}, \quad n_0 &= \cos \rho \cos z \in [-1, 0] , \quad z \in [-\pi, -\frac{\pi}{2}] \oplus [\frac{\pi}{2}, \pi] . \end{aligned} \quad (12.9a)$$

So one can use any of these two domains to cover the elliptical space model:

$$\begin{aligned} \tilde{G}^{(+1)}(\phi, z) : \quad \phi &\in [-\pi, +\pi] , \quad z \in [-\pi/2, +\pi/2] , \\ \tilde{G}^{(-1)}(\phi, z) : \quad \phi &\in [-\pi, +\pi] , \quad z \in [-\pi, -\pi/2] \oplus [+ \pi/2, +\pi] ; \end{aligned} \quad (12.9b)$$

The latter can be illustrated by the Fig 24:

Fig24 Domains $\tilde{G}^{(+1)}(\phi, z) - \tilde{G}^{(-1)}(\phi, z)$



It should be noted that the domain $\tilde{G}^{(-1)}$ consisting of two parts can be transformed to the domain $\tilde{G}^{(+1)}$ through the special change of variables:

$$\begin{aligned} z \in [\frac{\pi}{2}, \pi] , \quad z &= \pi - Z , \quad Z \in [0, +\frac{\pi}{2}] , \\ z \in [-\pi, -\frac{\pi}{2}] , \quad z &= -\pi - Z , \quad Z \in [-\frac{\pi}{2}, 0] , \end{aligned} \quad (12.10a)$$

Here the half-space $n_0^{(-1)} \in [-1, 0]$ is covered by

$$\begin{aligned} n_0 &= -\cos \rho \cos Z , \quad n_3 = \cos \rho \sin Z , \\ n_1 &= \sin \rho \cos \phi , \quad n_2 = \sin \rho \sin \phi , \end{aligned} \quad (12.10b)$$

in c^i -coordinates this looks as


$$c_1 = -\frac{\tan \rho}{\cos Z} \cos \phi , \quad c_2 = -\frac{\tan \rho}{\cos Z} \sin \phi , \quad c_3 = -\tan Z ; \quad (12.10c)$$

This referring of two domains

$$\tilde{G}^{(\delta=-1)}(\rho, \phi, z) \xrightarrow{z \rightarrow Z} \tilde{G}^{(\delta=+1)}(\rho, \phi, Z)$$

permits to establish identification rules for $\tilde{G}^{(\delta=-1)}(\rho, \phi, z)$. Indeed, from the scheme and relations


Fig 25 $\tilde{G}_1^{(\delta=+1)}(\phi, Z) : \rho \neq 0, \pi/2$



$$\begin{aligned} \tilde{G}_2(0, \phi, Z) : \quad & \rho = 0, \quad \vec{c} = -(0, 0, \tan Z), \\ \tilde{G}_3(0, \phi, Z) : \quad & \rho = \pi/2, \quad \vec{c} = -\infty (\cos \phi, \sin \phi, 0), \end{aligned} \quad (12.11a)$$

it follows the corresponding scheme and relations for $\tilde{G}_1^{(\delta=-1)}(\phi, z)$:

Fig 26 $\tilde{G}_1^{(\delta=-1)}(\phi, z) : \rho \neq 0, \pi/2$



$$\begin{aligned} \tilde{G}_2(0, \phi, z) : \quad & \rho = 0, \quad \vec{c} = (0, 0, \tan z), \\ \tilde{G}_3(0, \phi, z) : \quad & \rho = \pi/2, \quad \vec{c} = \infty (\cos \phi, \sin \phi, 0), \end{aligned} \quad (12.11b)$$

We will see below that two other variants combining previous ones $\delta = +1, -1$ may be of interest. In the Fig 24 they correspond to the domains (A), $z \in [0, +\pi]$ and (B), $z \in [-\pi, 0]$.

For definiteness, let us follow the variant $\delta = \pm 1$ and correspondingly two tetrads:

$$\begin{aligned} (\phi, z) \in \tilde{G}^{(+1)}(\phi, z), \quad & (\phi, z) \in \tilde{G}^{(-1)}(\phi, z) \\ dl^2(y^{(\delta)}) = d\rho^2 + \sin^2 \rho d\phi^2 + \cos^2 \rho dz^2, \\ e_{(i)j}^{(\delta)}(y) = \begin{vmatrix} 1 & 0 & 0 \\ 0 & \sin \rho & 0 \\ 0 & 0 & \cos \rho \end{vmatrix}. \end{aligned} \quad (12.12)$$

One should find an explicit form of the gauge matrix relating cartesian $e_{\beta}^{(\delta)(b)}(c)$ and cylindrical $e_{(a)}^{(\delta)\alpha}(y)$ tetrad bases in elliptical space.

$$L_a^{(\delta)b}(y) = e_{(a)}^{\alpha}(y) \frac{\partial c^{\beta}}{\partial y^{\alpha}} e_{\beta}^{(\delta)(b)}(c); \quad (12.13a)$$

With the use of (3.2), for $(\partial c^\beta / \partial y^\alpha)$ we have

$$\begin{aligned} \frac{\partial c^1}{\partial \rho} &= \frac{\cos \phi}{\cos^2 \rho \cos z}, & \frac{\partial c^2}{\partial \rho} &= \frac{\sin \phi}{\cos^2 \rho \cos z}, & \frac{\partial c^3}{\partial \rho} &= 0, \\ \frac{\partial c^1}{\partial \phi} &= -\frac{\tan \rho}{\cos z} \sin \phi, & \frac{\partial c^2}{\partial \phi} &= +\frac{\tan \rho}{\cos z} \cos \phi, & \frac{\partial c^3}{\partial \phi} &= 0, \\ \frac{\partial c^1}{\partial z} &= \frac{\tan \rho \sin z}{\cos^2 z} \cos \phi, & \frac{\partial c^2}{\partial z} &= \frac{\tan \rho \sin z}{\cos^2 z} \sin \phi, & \frac{\partial c^3}{\partial z} &= \frac{1}{\cos^2 z}, \end{aligned}$$

and the matrix $L_a^{(\delta) b}(y)$ reads (the notation is used: $(1 + \delta \cos \rho \cos z)^{-1} = \varphi$)

$$L_a^{(\delta) b}(\rho, \phi, z) = \begin{vmatrix} 1 & 0 \\ 0 & \delta \varphi (\cos \rho + \delta \cos z) \cos \phi \\ 0 & -\delta \sin \phi \\ 0 & \varphi \sin \rho \sin z \cos \phi \\ 0 & 0 \\ \delta \varphi (\cos \rho + \delta \cos z) \sin \phi & -\delta \varphi \sin \rho \sin z \\ \delta \cos \phi & 0 \\ \varphi \sin \rho \sin z \sin \phi & \delta \varphi (\cos z + \delta \cos \rho) \end{vmatrix}. \quad (12.13b)$$

By definition, the matrix $L_a^{(\delta=+1) b}(\rho, \phi, z)$ must be considered in the domain $\tilde{G}^{(+1)}(\rho, \phi, z)$, whereas the matrix $L_a^{(\delta=-1) b}(\rho, \phi, z)$ should be taken in $\tilde{G}^{(-1)}(\rho, \phi, z)$.

In isotropic form (for more detail see in [23]) the matrix $L^{(\delta)}$ is given by

$$U_{(\delta)}(\rho, \phi, z) = \frac{1}{2} \begin{vmatrix} (1 + \delta \varphi (\cos z + \delta \cos \rho)) & (1 - \delta \varphi (\cos z + \delta \cos \rho)) \\ (1 - \delta \varphi (\cos z + \delta \cos \rho)) & (1 + \delta \varphi (\cos z + \delta \cos \rho)) \\ -\delta \varphi \sin \rho \sin z & +\delta \varphi \sin \rho \sin z \\ -\delta \varphi \sin \rho \sin z & +\delta \varphi \sin \rho \sin z \\ \varphi \sin \rho \sin z e^{-i\phi} & \varphi \sin \rho \sin z e^{+i\phi} \\ -\varphi \sin \rho \sin z e^{-i\phi} & -\varphi \sin \rho \sin z e^{+i\phi} \\ \delta (1 + \delta \varphi (\cos z + \delta \cos \rho)) e^{-i\phi} & -\delta (1 - \delta \varphi (\cos z + \delta \cos \rho)) e^{+i\phi} \\ -\delta (1 - \delta \varphi (\cos z + \delta \cos \rho)) e^{-i\phi} & +\delta (1 + \delta \varphi (\cos z + \delta \cos \rho)) e^{+i\phi} \end{vmatrix} \quad (12.15a)$$

The isotropic matrix (12.15a) must be seen as parameterized in accordance with

$$U(\rho, \phi, z) = \begin{vmatrix} aa^* & cc^* & ac^* & a^*c \\ cc^* & aa^* & -ac^* & -a^*c \\ -ac & ac & aa & -cc \\ -a^*c^* & a^*c^* & -c^*c^* & a^*a^* \end{vmatrix}; \quad (12.15b)$$

where a, b stand for the elements of spinor matrix $B_{(\delta)}$:

$$B_{(\delta)}(\rho, \phi, z) = \begin{vmatrix} a & c \\ -c^* & a^* \end{vmatrix} = \begin{vmatrix} A(y) e^{i\alpha(y)} & C(y) e^{is(y)} \\ -C(y) e^{-is(y)} & A(y) e^{-i\alpha(y)} \end{vmatrix} \quad (12.15c)$$

As a first step, comparing (12.15a) and (12.15b), A and C are established

$$A^2 = \frac{1}{2}(1 + \delta \varphi (\cos z + \delta \cos \rho)) = \frac{(1 + \cos \rho)(1 + \delta \cos z)}{1 + \delta \cos \rho \cos z},$$

$$C^2 = \frac{1}{2}(1 - \delta \varphi (\cos z + \delta \cos \rho)) = \frac{(1 - \cos \rho)(1 - \delta \cos z)}{1 + \delta \cos \rho \cos z},$$

From the given A^2 and a^2 one finds $e^{-i\alpha}$:

$$\begin{aligned} e^{2i\alpha} A^2 &= a^2 = \delta e^{-i\phi} [1 + \delta \varphi (\cos z + \delta \cos \rho)] : \implies \\ e^{i\alpha} &= \sigma \sqrt{\delta} e^{-i\phi/2}, \quad e^{-i\alpha} = \sigma (\sqrt{\delta})^* e^{+i\phi/2}, \quad \sigma = \pm 1. \end{aligned}$$

Now, from given $-a^*c$, it follows expression for e^{is} :

$$\begin{aligned} -Ae^{-i\alpha}Ce^{is} &= -\varphi \sin \rho \sin z e^{+i\phi} \implies \\ e^{is} &= \frac{\varphi \sin \rho \sin z}{AC} e^{i\phi} e^{i\alpha} = \frac{\sin z}{\sqrt{\sin^2 z}} (\sigma \sqrt{\delta} e^{i\phi/2}). \end{aligned}$$

Thus, the spinor gauge matrix $B^{(\delta)}(\rho, \phi, z)$ is

$$\begin{aligned} A(y) &= +\sqrt{\frac{(1 + \cos \rho)(1 + \delta \cos z)}{2(1 + \delta \cos \rho \cos z)}}, \\ C(y) &= +\sqrt{\frac{(1 - \cos \rho)(1 - \delta \cos z)}{2(1 + \delta \cos \rho \cos z)}}, \\ e^{i\alpha(y)} &= \sigma \sqrt{\delta} e^{-i\phi/2}, \quad e^{-i\alpha(y)} = \sigma (\sqrt{\delta})^* e^{+i\phi/2}, \\ e^{is(y)} &= \sigma \sqrt{\delta} e^{i\phi/2} \frac{\sin z}{\sqrt{\sin^2 z}}, \quad e^{-is} = \sigma (\sqrt{\delta})^* e^{-i\phi/2} \frac{\sin z}{\sqrt{\sin^2 z}}, \end{aligned}$$

From (12.13a,b), taking into account identity

$$+\sqrt{\frac{1 - \cos z}{2}} \left(\frac{+\sqrt{\sin^2 z}}{\sin z} \right) = \sin \frac{z}{2},$$

for $B^{(+1)}(\rho, \phi, z)$ and $B^{(-1)}(\rho, \phi, z)$ we get

$$\begin{aligned} B_{(+1)}(\rho, \phi, z) &= \frac{1}{\sqrt{1 + \cos \rho \cos z}} \times \\ &\times \begin{vmatrix} \sqrt{1 + \cos \rho} \cos \frac{z}{2} e^{+i\phi/2} & \sqrt{1 - \cos \rho} \sin \frac{z}{2} e^{-i\phi/2} \\ -\sqrt{1 - \cos \rho} \sin \frac{z}{2} e^{+i\phi/2} & \sqrt{1 + \cos \rho} \cos \frac{z}{2} e^{-i\phi/2} \end{vmatrix}, \end{aligned} \quad (12.16a)$$

$$\begin{aligned} B_{(-1)}(\rho, \phi, z) &= \frac{1}{\sqrt{1 - \cos \rho \cos z}} \left(\frac{+\sqrt{\sin^2 z}}{\sin z} \right) \times \\ &\times \begin{vmatrix} \sqrt{1 + \cos \rho} \sin \frac{z}{2} i e^{+i\phi/2} & \sqrt{1 - \cos \rho} \cos \frac{z}{2} i e^{-i\phi/2} \\ \sqrt{1 - \cos \rho} \cos \frac{z}{2} i e^{+i\phi/2} & -\sqrt{1 + \cos \rho} \sin \frac{z}{2} i e^{-i\phi/2} \end{vmatrix}; \end{aligned} \quad (12.16b)$$

Below it will be used notation $+\sqrt{\sin^2 z}/\sin z = \text{Sgn } z \in \{+1, -1\}$.

By procedure used, the matrix $B_{(\delta=+1)}(\rho, \phi, z)$ should be considered in the domain $\tilde{G}^{(+1)}(\rho, \phi, z)$, and the matrix $B_{(\delta=-1)}(\rho, \phi, z)$ should be taken in the domain $\tilde{G}^{(-1)}(\rho, \phi, z)$. In the same time, one might employ them both in one the same domain, in such a context these two matrices would be connected by a local spinor gauge transformation associated with vector matrix (12.8).

In the vicinity of the point $(n_0 = +1, 0, 0, 0)$ (the limit of flat space) the matrix $B_{(+1)}$ will look

$$\rho \rightarrow 0, z \rightarrow 0: \quad B_{(+1)}(\rho, \phi, z) \rightarrow \begin{vmatrix} e^{+i\phi/2} & 0 \\ 0 & e^{-i\phi/2} \end{vmatrix}. \quad (12.17a)$$

As for the matrix $B_{(-1)}$, the vicinity of opposite point $(n_0 = -1, 0, 0, 0)$ should be taken:

$$\rho \rightarrow 0, \quad z = \pi - Z, \quad Z \rightarrow +0, \quad z = -\pi - Z, \quad Z \rightarrow -0$$

and

$$B_{(-1)}(\rho, \phi, z) \rightarrow \begin{vmatrix} i e^{+i\phi/2} & 0 \\ 0 & -i e^{-i\phi/2} \end{vmatrix}; \quad (12.17b)$$

13. On continuity condition for a fermion in elliptical space

Now we are ready to solve the main problem – analyzing the continuity condition for a fermion in elliptical space. There are two different bases playing the role of Cartesian in elliptical space:

$$\psi_{cart} = B^{-1}(\rho, \phi, z) \psi_{cyl}; \quad (13.1)$$

each variant with $\delta = +1$ and $\delta = -1$ presupposes its own domain $\tilde{G}^{(\delta)}(\rho, \phi, z)$ with respective identification rules on the boundary:

$(\delta = +1)$

$$\begin{aligned} \psi_1^{(\delta=+1)} &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos \rho \cos z}} \left[(1 + \cos \rho e^{-iz}) G_1(\rho) - i \sin \rho e^{-iz} G_2(\rho) \right], \\ \psi_2^{(\delta=+1)} &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos \rho \cos z}} \left[-i \sin \rho e^{+iz} G_1(\rho) + (1 + \cos \rho e^{+iz}) G_2(\rho) \right]. \end{aligned} \quad (13.2)$$

$(\delta = -1)$

$$\begin{aligned} \psi_1^{(\delta=-1)} &= (\text{Sgn } z) \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 - \cos \rho \cos z}} \left[(-1 + \cos \rho e^{-iz}) G_1(\rho) - i \sin \rho e^{-iz} G_2(\rho) \right], \\ \psi_2^{(\delta=-1)} &= (\text{Sgn } z) \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 - \cos \rho \cos z}} \left[-i \sin \rho e^{+iz} G_1(\rho) + (-1 + \cos \rho e^{+iz}) G_2(\rho) \right]. \end{aligned} \quad (13.3)$$


Because the requirement of continuity includes identity

$$\phi \implies \phi \pm \pi, \quad z \implies z \pm \pi.$$

it is easily seen that both variants, (13.2) and (13.3), lead us to the same result: no fermion solutions continuous in elliptical space exist.

One could try to construct continuous in $\tilde{S}_3$ solutions through the combination of variants (13.2) and (13.3). Let us examine this possibility in detail.

Variant (A):




Identification in horizontal boundaries is made in accordance with

$$\begin{aligned}
 (\phi, \quad z = +0) : \quad & c_1 = +\tan \rho \cos \phi, \quad c_2 = +\tan \rho \sin \phi, \quad c_3 = 0, \\
 (\phi + \pi, z = \pi - 0) : \quad & c_1 = +\tan \rho \cos \phi, \quad c_2 = +\tan \rho \sin \phi, \quad c_3 = 0; \quad (13.4a)
 \end{aligned}$$

the points of the internal dashed line in Fig 27 are associated with the infinite length vectors

$$\begin{aligned}
 (\phi, z = \pi/2 - 0) : \quad & c_1 = +\infty \tan \rho \cos \phi, \quad c_2 = +\infty \tan \rho \sin \phi, \quad c_3 = +\infty, \\
 (\phi, z = \pi/2 + 0) : \quad & c_1 = -\infty \tan \rho \cos \phi, \quad c_2 = -\infty \tan \rho \sin \phi, \quad c_3 = -\infty. \quad (13.4b)
 \end{aligned}$$

Variant (B):



Identification in horizontal boundaries is made in accordance with

$$\begin{aligned}
 (\phi, \quad z = -0) : \quad & c_1 = +\tan \rho \cos \phi, \quad c_2 = +\tan \rho \sin \phi, \quad c_3 = 0, \\
 (\phi + \pi, z = -\pi + 0) : \quad & c_1 = +\tan \rho \cos \phi, \quad c_2 = +\tan \rho \sin \phi, \quad c_3 = 0; \quad (13.5a)
 \end{aligned}$$

the points of the internal dashed line in Fig 27 are associated with the infinite length vectors

$$(\phi, z = -\pi/2 + 0) : \quad c_1 = +\infty \tan \rho \cos \phi, \quad c_2 = +\infty \tan \rho \sin \phi, \quad c_3 = -\infty,$$

$$(\phi, z = -\pi/2 - 0) : \quad c_1 = -\infty \tan \rho \cos \phi, \quad c_2 = -\infty \tan \rho \sin \phi, \quad c_3 = +\infty, \quad (13.5b)$$

The Variants (A) and (B) lead to the fermion functions of the form: Variant (A)

$$\begin{aligned} z \in [0, +\frac{\pi}{2}] : \quad \psi_1^{(\delta=+1)} &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos \rho \cos z}} [(1 + \cos \rho e^{-iz})G_1 - i \sin \rho e^{-iz}G_2], \\ z \in [\frac{\pi}{2}, +\pi] : \quad \psi_1^{(\delta=-1)} &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 - \cos \rho \cos z}} [(1 - \cos \rho e^{-iz})G_1 + i \sin \rho e^{-iz}G_2]; \\ z \in [0, +\frac{\pi}{2}] : \quad \psi_2^{(\delta=+1)} &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos \rho \cos z}} [-i \sin \rho e^{+iz}G_1 + (1 + \cos \rho e^{+iz})G_2], \\ z \in [\frac{\pi}{2}, +\pi] : \quad \psi_2^{(\delta=-1)} &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 - \cos \rho \cos z}} [+i \sin \rho e^{+iz}G_1 + (1 - \cos \rho e^{+iz})G_2]. \end{aligned} \quad (13.6)$$

Variant (B)

$$\begin{aligned} z \in [0, -\frac{\pi}{2}] : \quad \psi_1^{(\delta=+1)} &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 + \cos \rho \cos z}} [(1 + \cos \rho e^{-iz})G_1 - i \sin \rho e^{-iz}G_2], \\ z \in [-\pi, -\frac{\pi}{2}] : \quad \psi_1^{(\delta=-1)} &= \frac{e^{i(m-1/2)\phi} e^{i(k+1/2)z}}{\sqrt{1 - \cos \rho \cos z}} [(-1 + \cos \rho e^{-iz})G_1 - i \sin \rho e^{-iz}G_2], \\ z \in [0, -\frac{\pi}{2}] : \quad \psi_2^{(\delta=+1)} &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 + \cos \rho \cos z}} [-i \sin \rho e^{+iz}G_1 + (1 + \cos \rho e^{+iz})G_2], \\ z \in [-\pi, -\frac{\pi}{2}] : \quad \psi_2^{(\delta=-1)} &= \frac{e^{i(m+1/2)\phi} e^{i(k-1/2)z}}{\sqrt{1 - \cos \rho \cos z}} [-i \sin \rho e^{+iz}G_1 + (-1 + \cos \rho e^{+iz})G_2], \end{aligned} \quad (13.7)$$

It is easily seen that both in (A) and (B), one can satisfy $\psi(M) = \psi(M')$, where M and M' belong to vertical and horizontal boundaries, however such solutions would take different values in the identified point $\pm\infty\vec{c}_0$ of the internal dashed lines. Therefore, the variants (A) and (B) give the same result: any Dirac solutions continuous in elliptical space model do not exist.

Supplement A. Cartesian and cylindrical tetrads in S_3 .

In S_3

$$n_0^2 + n_1^2 + n_2^2 + n_3^2 = \rho^2, \quad (A.1)$$

one may introduce (quasi) cartesian coordinates and tetrad

$$x^1 = \frac{n_1}{1 + n_4}, \quad x^2 = \frac{n_2}{1 + n_4}, \quad x^3 = \frac{n_3}{1 + n_4}; \quad (A.2a)$$

$$n_i = \frac{2x^i}{1 + x^2}, \quad n_4 = \frac{1 - x^2}{1 + x^2}, \quad , \quad x^2 = x^1x^1 + x^2x^2 + x^3x^3; \quad (A.2b)$$

$$dl^2(x) = \frac{1}{f^2} [(dx^1)^2 + (dx^2)^2 + (dx^3)^2] , \quad f = \frac{1+x^2}{2} ; \quad (A.2c)$$

$$e_{(a)}^\alpha(x) = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & 1/f & 0 & 0 \\ 0 & 0 & 1/f & 0 \\ 0 & 0 & 0 & 1/f \end{vmatrix} ; \quad (A.2d)$$

here a and α take values $0, 1, 2, 3$; (a) stands for a tetrad index; α stands for a generally covariant index. Also, in the spherical space one can introduce cylindrical coordinates and tetrad:

$$n_1 = \sin \rho \cos \phi , \quad n_2 = \sin \rho \sin \phi , \quad n_3 = \cos \rho \sin z , \quad n_4 = \cos \rho \cos z ,$$

$$dl^2(y) = (d\rho^2 + \sin^2 \rho d\phi^2 + \cos^2 \rho dz^2) , \quad e_{(a)}^\alpha(y) = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \sin^{-1} \rho & 0 \\ 0 & 0 & 0 & \cos^{-1} \rho \end{vmatrix} ; \quad (A.3)$$

$$G(\rho, \phi, z) = \{ \rho \in [0, +\pi/2] , \quad \phi \in [-\pi, +\pi] , \quad z \in [-\pi, +\pi] \} .$$

It is the matter of simple calculation to find a vector and spinor gauge matrices relating Cartesian and cylindrical bases in space S_3 . The vector matrix obeys the equation

$$e_{(a)}^\alpha(y) = \frac{\partial y^\alpha}{\partial x^\beta} L_a^b(y) e_{(b)}^\beta(x) ; \quad (A.4a)$$

from this it follows

$$L_a^b(y) = e_{(a)}^\alpha(y) \frac{\partial x^\beta}{\partial y^\alpha} e_{(b)}^\beta(x) . \quad (A.4b)$$

With the use of the formulas

$$x^1 = \frac{\sin \rho \cos \phi}{1 + \cos \rho \cos z} , \quad x^2 = \frac{\sin \rho \sin \phi}{1 + \cos \rho \cos z} , \quad x^3 = \frac{\cos \rho \sin z}{1 + \cos \rho \cos z} ; \quad (A.5)$$

$$\frac{\partial x^1}{\partial \rho} = \frac{(\cos \rho + \cos z) \cos \phi}{(1 + \cos \rho \cos z)^2} , \quad \frac{\partial x^2}{\partial \rho} = \frac{(\cos \rho + \cos z) \sin \phi}{(1 + \cos \rho \cos z)^2} , \quad \frac{\partial x^3}{\partial \rho} = - \frac{\sin \rho \sin z}{(1 + \cos \rho \cos z)^2} ,$$

$$\frac{\partial x^1}{\partial \phi} = - \frac{\sin \rho \sin \phi}{(1 + \cos \rho \cos z)} , \quad \frac{\partial x^2}{\partial \phi} = \frac{\sin \rho \cos \phi}{(1 + \cos \rho \cos z)} , \quad \frac{\partial x^3}{\partial \phi} = 0 ,$$

$$\frac{\partial x^1}{\partial z} = \frac{\sin \rho \cos \rho \sin z}{(1 + \cos \rho \cos z)^2} \cos \phi , \quad \frac{\partial x^2}{\partial z} = \frac{\sin \rho \cos \rho \sin z}{(1 + \cos \rho \cos z)^2} \sin \phi , \quad \frac{\partial x^3}{\partial z} = \frac{(\cos \rho + \cos z) \cos \rho}{(1 + \cos \rho \cos z)^2} .$$

one can determine the explicit form of $L_a^b(y)$ (below $\varphi = (1 + \cos \rho \cos z)^{-1}$)

$$L_a^b(y) = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & \varphi(\cos \rho + \cos z) \cos \phi & \varphi(\cos \rho + \cos z) \sin \phi & -\varphi \sin \rho \sin z \\ 0 & -\sin \phi & \cos \phi & 0 \\ 0 & \varphi \sin \rho \sin z \cos \phi & \varphi \sin \rho \sin z \sin \phi & \varphi(\cos \rho + \cos z) \end{vmatrix} . \quad (A.6)$$

By the given vector matrix $L_a^b(\rho, \phi, z)$ one should determine a corresponding spinor matrix

$$B(\rho, \phi, z) = \begin{vmatrix} a & c \\ -c^* & a^* \end{vmatrix} , \quad B \in SU(2) , \quad a a^* + c c^* = +1 .$$

It is convenient to employ isotropic representation for the above vector matrix $L(y) \implies U(y) = S L(y) S^{-1}$:

$$S = \frac{1}{\sqrt{2}} \begin{vmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & -1 \\ 0 & 1 & -i & 0 \\ 0 & 1 & +i & 0 \end{vmatrix}, \quad S^{-1} = \frac{1}{\sqrt{2}} \begin{vmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & i & -i \\ 1 & -1 & 0 & 0 \end{vmatrix}. \quad (A.7)$$

In isotropic basis the Lorentz matrix must be parameterized in accordance with ⁶

$$U(\rho, \phi, z) = \begin{vmatrix} aa^* & cc^* & ac^* & a^*c \\ cc^* & aa^* & -ac^* & -a^*c \\ -ac & ac & aa & -cc \\ -a^*c^* & a^*c^* & -c^*c^* & a^*a^* \end{vmatrix}. \quad (A.8)$$

In this way for the spinor matrix one find the expression (it suffices to follow the 2-component spinor)

$$\xi_{cyl} = B(\rho, \phi, z) \xi_{cart} : \quad B(\rho, \phi, z) = \sigma \begin{vmatrix} A e^{+i\phi/2} & C e^{-i\phi/2} \\ -C e^{+i\phi/2} & A e^{-i\phi/2} \end{vmatrix}, \quad (A.9)$$

$$A = \sqrt{\frac{1 + \cos \rho}{1 + \cos \rho \cos z}} \cos \frac{z}{2}, \quad C = \sqrt{\frac{1 - \cos \rho}{1 + \cos \rho \cos z}} \sin \frac{z}{2}.$$

More details on the properties of that spinor transformation see in [23].

⁶For more detail see [23].

REFERENCES

1. Peter F., Weyl H. Die Vollständigkeit der primitiven Darstellungen einer geschlossenen kontinuierlichen Gruppe. // Math. Ann. 1927. Bd. 97. S. 737-755. Russian translation in: On completeness of primitive representations of a compact continuous group. Yspexi matem. nauk. 1936. Tome 2. 144-160; Weyl H. Harmonics on homogeneous manifolds. Math. Ann. 1934. Vol. 35. P. 486-499.
2. Barut A., Ronchka R. Theory of the representations on the groups ant its appllications. Vol. 1. 1980 (in Russian).
3. Tretyakova N.N. Harmonic analysis in homogeneous spaces. Minsk. 1996 (in Russian).
3. Wigner E E. Group theory and its applications to the quantum theory of atomic spectra. New York. 1959; Moscow. 1961 (in Russian)
4. Fano U., Racah G. Irreducible tensorial sets. New York. 1959.
5. Edmonds A. Angular momentums in quantum mechanics. Prinston. 1957; See in: Deformation of atomic nucleuses. Moscow. 1958. Pages 305-351 (in Russian);
6. Brink D.M., Satchler G.R. Angular momentum. Oxford. 1962.
12. Bopp F., Haag. Z. Naturf. 1950. Bd. 5a. S. 644.
11. Gelfand I.M., Shapiro E.Ya. Represenrtations of the rotation group of the 3-dimensional space and its application. Uspexi matem. nauk. 1952. Tome 7. Pages 2-... (in Russian).
11. Godement R. A theory of spherical functions. Trans. Amer. math. Soc. 1952. Vol. 73. pages 496-556.
14. Kahan T. Theorie des groupes en physique classique et quantique. Tome 1. Paris, 1960.
16. Lyubarski G.Ya. The application of the group theory in physics. Oxford. 1960.
15. Yutsis A.P., Levinson I.B., Vanagas V.V. The theory of angular momentum. Vilnus. 1960 (in Russian).
13. Schwinger J. Quantum theory of angular momentum. ed. by L.C. Biedenharn and H. van Dam. new York. 1965.
16. Vilenkin N.Ya. Special function and the the theory of representations of the groups. Moscow. 1965(in Russian).
17. Varshalovich D.A., Moskalev A.N., Xersonskiy V.K. Quantum theory of angular momentum. Leningrad. 1975 (in Russian).
20. Biedenharn L.C., Lauck J.D. Angular momentum in quantum physics. Volumes 1-2. Moscow. 1964 (in Russian).
21. Junker J. E., De Vries E. A note on orthogonality and completness of the rotation matrices. Nucl. Phys. 1967. Vol A105. P. 621-626.
21. Taylor M.E. Fourier series on compact Lie groups. Proc. Amer. Math. Soc. 1968. Vol. 19. P. 110301105.
7. Davydov A.S. Quantum mechanics. Moscow. 1973 (in Russian).
8. Landau L.D., Lifshits E.M. Quantum mechanics, non-relativistic theory. Moxcow. 1974 (in Rissian).
23. Lee Beers, Richard S. Millman. Analytic vector harmonic expansions on $SU(2)$ and S^2 . J. math. Phys. 1975. Vol. 16. P. 11-19.
23. Vernon D. Sandberg, Tensor spherical harmonics on S^2 and S^3 as eigenvalue problem. J. Math. Phys. 1978. Vol 19. P. 2441-2446.
9. Gibson W., Pollard B. Symmetry principles in particle physics. Moscow. 1979)in Russian).
22. Madelung E. Mathematical apparatus in physics. Moscow. 1968 (in Russian).
10. Fedorov F.I. The Lorentz group. Moscow. 1979 (in Russian).
- 23.

23. Olevskiy M.N. Three-orthogonal systems in spaces of constant curvature in which equation $\Delta_2 U + \lambda U = 0$ permits the full separation of variables. Matem. Sbornik. 1950 P. 379-426.
23. Red'kov V.M. Gauge transformations for a fermion function in spherical space. Riemanian manifold of constant positive curvature with spinor structure. Preprint 669. Institute of Physics, Academy of Sciences of Belarus. Minsk. 1993. 35 pages.
24. Red'kov V.M. On spinor structure of the pseudo Riemanian space-time and property of global continuity for fermionic wave functions. //Vesti AN BSSR. ser. fiz.-mat. nauk. 1994. 3. Pages 49-55.